



PREVALENCE ANALYSIS INCIDENCE OF UPPER RESPIRATORY COVID-19 - INFECTED PATIENTS

1. **Mirzoyeva Mexriniso Rizoyevna**

2. **Sadullayeva Rushana
Sadullayevna**

Received 29th Nov 2023,
Accepted 29th Dec 2023,
Online 31th Jan 2024

^{1,2} Bukhara State Medical Institute

Abstract: The material of the study was 25 patients for 2019-2021 up to 45 years old with COVID-19 infection. 10 patients were examined in the emergency department and 15 patients in the hospital. Children were examined regardless of the presence of complaints. In addition to standard research methods (general blood count, urine, bacteriological and biochemical studies), we conducted a thorough otorhinolaryngological examination in all patients, and in 2 (8%) X-ray examinations.

Key words: patients, COVID-19, upper respiratory tract disease.

Recently, there has been a global pandemic burden due to human-to-human transmission of a new coronavirus disease like COVID-19. Since the outbreak in December 2019, COVID-19 has affected more than a dozen million people [2,4]. As is known, the majority of COVID-19 cases among people are mild (80%), 20% of infected patients may develop a serious illness, and 5% may develop severe illness and pneumonia, or acute respiratory distress syndrome, which requires artificial ventilation and hospitalization in the intensive care unit [8,10]. Lesions of the mucous membrane of the oral cavity and upper respiratory tract demonstrated various clinical aspects, differing in localization, size, color and quantity. Blisters, ulcers, erosions, spots and plaques were observed in patients [1, 3, 11].

Current studies show that the coronavirus enters human cells through the angiotensin converting enzyme receptor 2 (ACE2), during the studies, organs at risk and vulnerable to

infection with coronavirus 2 (SARS-CoV-2) of severe acute respiratory syndrome were identified [4,6]. Thus, cells with the distribution of ACE2 receptors can become host cells for the virus and cause an inflammatory reaction in related organs and tissues, such as the mucous membrane of the tongue and salivary glands [5,9]. The interaction of SARS-CoV-2 with ACE2 receptors can also impair the sensitivity of taste receptors, which can cause dysfunctional taste reactions.

History shows that in December 2019, the first cases of pneumonia caused by an unknown infectious agent were reported in Wuhan, China. It was later discovered that the infectious agent is a new beta-coronavirus (β CoV), which was named coronavirus 2 of severe acute respiratory syndrome (SARS-CoV-2) because of its phylogenetic similarity to SARS-CoV. Infections soon became widespread in China and the rest of the world,

which led to the WHO declaring a pandemic and a "public health emergency of international importance" [7.10-14].

The so-called coronavirus disease 2019 (COVID-19) has a higher severity and mortality in the elderly, in patients with concomitant diseases such as hypertension and diabetes, as well as in people with reduced immune activity. Apart from the fact that the pandemic poses a serious threat to public health, it has also affected social life and the global economy.

The aim of the study was the epidemiological characteristics of upper respiratory tract diseases in COVID-19 infected patients.

Materials and methods of research.

During the period from May 2019 to July 2021, 25 patients with a diagnosis of COVID-19 infection were under our supervision. The research was conducted on the basis of the regional Children's Infectious diseases hospital. In (32%) 8 patients with COVID-19, the status was established upon admission to the hospital. The material of the study was 35 patients for 2019-2021 under 45 years of age with COVID-19 infection. 10 sick children were examined in the reception departments and 15 patients in the hospital. Patients were examined regardless of the presence of complaints. In addition to standard methods of examination (general analysis of blood, urine, bacteriological and biochemical studies), we conducted a thorough otorhinolaryngological examination for all children, and X-ray examination in 2 (8%).

Careful observation of the patient's behavior and subsequent analysis of the semiotics of the disease revealed; ear pain, (16%) pus from the ear (32%), anxiety of the patient (40%), poor sleep (100%), pain when swallowing (32%), nasal congestion (18%), discharge from nose (18%) sore throat (8%) cough (8%). We believe that a doctor should pay attention to weakness at high body temperature in a patient, as it may be a consequence of depression of the nervous system. We observed this symptom in the clinic of acute otitis media in patients only in 5 patients (20%).

According to age, the patients are divided into 4 groups:

1st group from 18 to 25 years old;

2nd group from 26 years to 30 years;

3rd group from 31 years to 40 years;

4th group from 41 to 45 years old;

The incidence of acute otitis media in men is slightly higher (25%), compared with women (20.4%) at the same time, catarrhal otitis media is twice as common as acute purulent otitis media (30.9% and 14.5%), respectively. The incidence of tonsillitis and rhinosinusitis in the 1st group of men is slightly higher (16%), compared to women (8% and 12%), at the same time acute laryngitis occurs in men (8%) and women (12%), respectively, acute pharyngitis was not observed in this group. In the second group, the incidence of acute otitis media in men and women was almost the same, but acute catarrhal otitis media occurs almost 5 times more often than acute purulent otitis media 31.3% and 6.2%, respectively. The incidence of acute tonsillitis in men and women was almost the same (12% and 12%, respectively) acute laryngitis and pharyngitis were not observed in this group.

Rhinosinusitis occurs in men (8%), and in women 16%, respectively. The incidence of acute otitis media in the 3rd group of patients was the highest (46.3%), men were sick more often than women (28.2% and 18.1%), respectively.

The incidence of acute tonsillitis in this group was more common for men (16%) than for women (16% and 8%), respectively. Acute rhinosinusitis affected almost the same (12% and 12%), respectively, acute pharyngitis affected men (12%) and women (8%).

In group 4, the incidence of acute otitis media was also higher in men compared to women. ((17.6%-10.2%), and tonsillitis occurs in men (12%), in women (16%), rhinosinusitis in men (16%), girls (8%), pharyngitis occurs in boys (8%), in women (12%), respectively.

We also compared the results of ENT pathology detection in the emergency department and in the hospital. The frequency of ENT pathology registrations in the emergency department was significantly lower than that in the hospital, although all patients examined in the hospital had already passed through the emergency department. This is primarily due to the absence of a permanent ENT doctor in the emergency department, where an infectious disease specialist forced to take many patients a day does not always pay attention to ENT pathology or pathology of the upper respiratory tract.

Thus, the incidence of ENT organs in COVID-19-infected patients averages 70-80%, at the same time, due attention is not paid in the emergency department to the presence of ENT pathology, adequate therapy is not prescribed, which contributes to the development of a chronic condition or various intracranial complications, which ultimately can lead to disability of varying degrees.

Conclusions:

1. Acute inflammation of the mucous membrane of the upper respiratory tract in COVID-19-infected patients occurs from 70% to 80% of cases.
2. In infected hospitals, it is necessary to allocate the staff of an otorhinolaryngologist for mandatory examination of all COVID-19-infected incoming patients in order to prevent possible secondary development of intracranial complications.

Literature

1. Khamidova, N. K., Mirzoeva, M. R., & Narzullaev, N. U. (2021). Clinical and Immunological Study of the Effect of Different types of Therapy on the Course of Allergic Rhinitis in Children with Hymenolepiasis. *Annals of the Romanian Society for Cell Biology*, 1900-1908.
2. Keldiyorova, Z. D., Ostonova, G. S., Mirzoeva, M. R., & Narzullaev, N. U. (2021). State of the immune system in children with infectious mononucleosis. *New day in medicine. Бухоро-1* (33), 283-286.
3. Mirzoeva, M. R., & Khamidova, N. K. (2020). Clinical and epidemiological characteristic of hymenolepidosis children. *AcademiciaAn International Multidisciplinary Research Journal*, 169-173.
4. Мирзоева, М. Р., Нарзуллаев, Н. У., & Хамидова, Н. К. (2020). Клинико-эпидемиологическая характеристика гименилипедоза у детей. *Новый день в медицине*, (2), 441-443.
5. Keldiyorova, Z. D., Narzullaev, N. U., & Mirzoeva, M. R. (2022). Immunological disorders in infectious mononucleosis in children. *Neuroquantology*, 20(6), 9600.
6. Келдиёрова, З. Д., Нарзуллаев, Н. У., & Мирзоева, М. Р. (2020). Цитокиновый профиль детей с острым воспалением небного миндалина при острой инфекционной мононуклеозе на фоне лечения. *Тиббиётда янги кун*, (2), 30.
7. Мирзоева, М., Бабаходжаев, С., Халилова, З., & Саъдинов, П. (2015). Характер и частота осложнений у больных хроническим токсоплазмозом. *Журнал проблемы биологии и медицины*, (1 (82)), 52-54.
8. Мирзоева, М. Р., Худойдодова, С. Г., & Фарманова, М. А. (2019). Кишечные паразитозы у детей с гипопигментозом кожи и их возможная связь с этиопатогенезом заболевания. *Сопредседатели редакционной коллегии: АШ ИНОЯТОВ*, 57.
9. Mirzoyeva, M. R., Ostonova, G. S., & Narzullaev, N. U. (2021). The advancement of nonspecific prophylaxis and therapy pseudotuberculosis and intestinal versiniosis.

10. Mirzoeva, M. R., & Ostonova, G. S. (2021). In the modern classification, the place of intestinal yersiniosis. *Innovative Technologica Methodical Research Journal*, 2(09), 8-12.
11. Мирзоева, М. Р. (2018). Характер и частота осложнений у больных хроническим токсоплазмозом. *Редакционная коллегия*, 207.
12. Мирзоева, М. Р., Бабаходжаев, С. Н., & Сайфутдинова, Н. Н. (2015). Характер лабораторных изменений у больных хроническим токсоплазмозом. *Журнал теоретической и клинической медицины*, (2), 124-125.
13. Мирзаева, М. А., & Адилханова, А. Н. (2014). Формирование микрофлоры кишечника у новорожденных в экологически неблагоприятном регионе республики Узбекистан. *Проблемы биологии и медицины*, (3), 79.
14. Mirzoeva, M. R., & Mavlyanova, S. Z. роль гелминтно-паразитарной сенсibilизации в клиническом течении PITYRIASIS ALBA. VA *ESTETIK TIBBIYOT*, 125.

