

Article

# Preparation And Evaluation Of The Biological Activity Of A 2-Amino Pyran Ring Using A Solid Base Catalyst

Mohammed Jwher Saleh<sup>1</sup>, Jamil Nadhem Saleh<sup>2</sup>, Khalid al-badrany<sup>3</sup>, Adil Hussein Dalaf<sup>4</sup>, Reem suhail Najm<sup>5</sup> Abdul Wahed Abdul Sattar Talluh<sup>6</sup>

1. Salah al-Din Education Directorate/Iraq
- \* Correspondence: [jamelnadhemsaleh@gmail.com](mailto:jamelnadhemsaleh@gmail.com)
2. Salah al-Din Education Directorate/Iraq
3. Tikrit University, College of Education for Pure Sciences/Departement of Chemistry
4. Department of Chemistry, College of Science, Tikrit University, Tikrit-Iraq
5. Department of Physiology, Pharmacology and Biochemistry, College of Veterinary Medicine, University of Tikrit, Salahuddin-Tikrit, Iraq
6. Tikrit University, College of Basic Education, Shirqat, Tikrit, Iraq

**Abstract:** The research included preparing a basic catalyst by reacting potassium nitrate with alumina to produce the heterogeneous solid basic catalyst and then introducing the catalyst as a catalyst in preparing the 2-amino pyran ring by reacting the prepared chocolate with malononitrile and using only stirring without sublimation or heating. The compositions were validated using physical measurements such as percentage, melting point, and colour and spectroscopic measurements such as proton and carbon nuclear magnetic resonance and infrared spectra. The nanoscale size of the catalyst was also confirmed by scanning electron microscope (SEM) analysis. Two types of positive bacteria were used – gram and gram-negative to verify biological activity.

**Keywords:** catalyst, pyran, biological activity.

**Citation:** Mohammed Jwher Saleh. Preparation And Evaluation Of The Biological Activity Of A 2-Amino Pyran Ring Using A Solid Base Catalyst. Central Asian Journal of Medical and Natural Science 2024, 5(4),130-138.


Received: 10<sup>th</sup> Apr 2024  
Revised: 11<sup>th</sup> Mei 2024  
Accepted: 24<sup>th</sup> Jun 2024  
Published: 25<sup>th</sup> Jul 2024



**Copyright:** © 2024 by the authors. Submitted for open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>)

## 1. Introduction

**Catalyst** The term catalyst was originally defined by the scientist Berzelius in 1836 as a chemical substance added in small quantities to the reacting medium to increase the rate of a chemical reaction by reducing the energy needed to reach the activated complex phase in which the reactants move through their molecular orbitals and then decompose to produce the products. The catalyst does not cause thermodynamically impossible reactions but rather accelerates possible reactions. It has been found that adding small amounts of metallic materials, such as transition elements, to the centre of some reactions increases reaction rates compared to the absence of these materials [1]. Pyran is a heterocyclic, unsaturated (non-aromatic) hexacyclic compound consisting of six main atoms, five of which are carbon atoms and one oxygen atom [2], and has the chemical formula C<sub>5</sub>H<sub>6</sub>O. Its systematic names (IUPAC) are 2H-Pyran, and 4H-Pyran, and there are other names: H-Oxine, 4H-Oxine<sup>2</sup>. There are isomers of pyran, with different positions of the double bond. In 2H-pyran, the saturated carbon is at the 2-position, while in 4H-pyran, the saturated carbon is at the 4-position [3].



They have a variety of fascinating biological functions and significant significance in organic biochemistry [4]. The presence of 4H-cyanohydrins and their derivatives in a large variety of physiologically active chemicals makes them a significant class of molecules. Applications of 4H-cyanopyran in modern pharmaceutical chemistry include antioxidant and antibacterial properties [5], anticoagulant, antiallergic [8], antituberculosis [6], and antitumor [7].

## 2. Materials and Methods

**2.1 Chemical Used,** Chemicals prepared by Aldrich, BDH Thomas, Fluka, and Merck were used

**2.2 Instruments Used,** The melting point is measured using a thermometer 9300, KBr disk with a scale of 400-4000  $\text{cm}^{-1}$ , FT-IR 8400S Shimadzu spectrophotometer,  $^1\text{H}$ - and  $^{13}\text{C}$ -NMR spectra of Bruker equipment operating at 400 MHz. Thin-layer chromatography (TLC) was analysed using 0.2 mm thick Fluka silica gel plates.

**2.3 Preparation of catalyst derivatives.** [9]

Weigh (1 mol) potassium nitrate and (3 mol) alumina and grind them evenly in a mortar, then add 5 drops of deionized water during the grinding process until uniform, form a paste, and dry at 100 °C. It is activated at a temperature of 110 degrees Celsius for an hour, then the mixture is placed in a ceramic bowl and activated at a temperature of 600 degrees Celsius for 3 hours. The catalyst was left to cool in the air.

**2.3 Preparation of 2-amino pyran derivatives (M36-M40).**[10,11]

Dissolve (0.015 mol) chalcone in (15 ml) ethanol in a circular flask (15 ml) equipped with a magnetic stirrer, then add (0.225 mol, 0.148 ml) malononitrile and stir for (15) minutes (11), and add the catalyst ( $\text{Al}_2\text{O}_3\text{-OK}$ , 20% wt chalcone) Stir in a water bath at (40)°C for 3-4 hours. The solution was then filtered, the precipitate was discarded, and the filtrate was allowed to dry and recrystallized from absolute ethanol. Use TLC plates to track the reaction process. Table (2-10) shows in Table (MH36-MH40).


Comp No.	R	Molecular formula	m.p. °C	Yield %	Colour
MH <sub>16</sub>	4-Cl	$\text{C}_{19}\text{H}_{12}\text{BrClN}_3$	165-167	42	Off white
MH <sub>17</sub>	4-NO <sub>2</sub>	$\text{C}_{19}\text{H}_{12}\text{BrN}_4\text{O}_2$	183-185	38	Brown
MH <sub>18</sub>	4-OCH <sub>3</sub>	$\text{C}_{20}\text{H}_{15}\text{BrN}_3\text{O}$	116-118	40	Blue
MH <sub>19</sub>	4-F	$\text{C}_{19}\text{H}_{12}\text{BrFN}_3$	207-209	46	Light green
MH <sub>20</sub>	2,3-diCl	$\text{C}_{19}\text{H}_{11}\text{BrCl}_2\text{N}_3$	143-145	57	Orange

## 2.4 Biological activity study

*Staphylococcus aureus* is both gram-negative and gram-positive. The pathogen used in this study is *Escherichia coli*. The Department of Pure Science and Life Sciences Education uses Molton-Hinton agar as a bacterial growth medium. Chemical solutions of M36, M37, M39, and M40 were prepared using dimethyl sulfoxide (DMSO) at concentrations of (0.01, 0.001, 0.0001) mg/mL [12, 13]. Determine and monitor the minimum inhibitory concentration (MIC). Mueller Hinton agar was used as a nutrient medium and diffusion techniques were performed to confirm the susceptibility of the bacterial isolates used in the study. After preparing the culture medium, sterilize it, distribute it onto Petri dishes, and allow it to freeze. Next, drill four small holes in each plate. They were then incubated at 37°C for a full day. Derivatives used. These indicate the sensitivity of the derivatives used. As the diameter increases, these derivatives depend on the damping diameter of the plate surrounding the hole used. When the chemical produced shows an inhibitory effect, its biological activity increases, similar to the inhibitory effects of antibiotics. [14,15].

## 3. Results and Discussion


2-Aminopyran compounds were prepared according to the following scheme



**Scheme (1): Path of the Ready Compounds (M36-M40)**


### 3.1 Characterization of catalyst

The proposed mechanism for preparing the catalyst was as follows:



**Scheme; 2 Mechanism of catalytic reaction**


SEM analysis of the catalyst shown in Figure 1 shows that the cross-sectional area (2  $\mu\text{m}$ ) and peak radius of the catalyst particles is 2.529 nm belonging to nanocrystals [16].



**Figure (1):** shows an SEM image of the catalyst

### 3.2 Characterization of 2- Amino pyran derivatives (M36-M40)

The proposed mechanism for preparing the 2-AMINO PYRAN was as follows:



**Scheme (3):** Mechanism of preparation of 2-aminopyran derivatives [MH36- MH40]

Examining the 2-aminopyran derivative's infrared spectrum (FT-IR) revealed two absorption bands that belonged to the bond (NH<sub>2</sub>) and appeared in the interior in the ranges of (3240-3311) cm<sup>-1</sup> and (3320-3375) cm<sup>-1</sup>. Additionally, an absorption band in the range of (3047-3078) cm<sup>-1</sup> also appeared as a result of the aromatic bond's expansion. Additionally, it was noted that absorption bands appeared of (2168-2216) cm<sup>-1</sup> due to the expansion of aromatic bonds (CN) and (1583-1595) cm<sup>-1</sup> due to the expansion of the alkene affinity band (C=C); an absorption band appears in the range of (1360-1390) cm<sup>-1</sup> due to the stretching of the affinity group (CO-O); and two absorption bands appear in the ranges of (2937-2962) cm<sup>-1</sup> and (2824-2910) cm<sup>-1</sup> that is associated with the stretching of the aliphatic bond (C-H) as demonstrated [16]. Table (3-9).

Table (2): Infrared absorption results (cm<sup>-1</sup>) for [MH36-MH40] compounds

Comp. No.	R	IR (KBr) cm <sup>-1</sup>						
		$\nu(\text{C-H})$ Arm	$\nu(\text{C-O})$	$\nu(\text{C-H})$ .Aliph.	$\nu(\text{NH}_2)$	$\nu(\text{C=C})$	$\nu(\text{C=C})$ Arom.	Others
MH <sub>36</sub>	4-Cl	3060	1390	2945 2844	3371 3311	1589	1512 1463	$\nu(\text{C-Br})$ 622 $\nu(\text{CN})$ 2179
MH <sub>37</sub>	4-NO <sub>2</sub>	3078	1377	2937 2824	3345 3240	1588	1519 1454	$\nu(\text{C-Br})$ 654 $\nu(\text{CN})$ 2168
MH <sub>38</sub>	4-OCH <sub>3</sub>	3047	1360	2939 2841	3325 3260	1595	1515 1490	$\nu(\text{CBr})$ 609 $\nu(\text{-CN})$ 2196
MH <sub>39</sub>	4-F	3051	1384	2950 2896	3375 3303	1583	1514 1463	$\nu(\text{C-Br})$ 667 $\nu(\text{CN})$ 2216
MH <sub>40</sub>	2,3di-Cl	3054	1385	3962 3910	3320 3244	1593	1537 1487	$\nu(\text{C-Br})$ 619 $\nu(\text{CN})$ 2184




Figure (2): The infrared spectrum of the compound [MH36]




Figure (3): The infrared spectrum of the compound [MH39]

The  $^1\text{H}$ -NMR spectrum of the compound [MH36] shows that multiple signals attributed to the aromatic ring protons appear in the ppm range (7.18-7.91), and a signal at the ppm position (4.06) attributable to the aromatic ring protons that is consistent with the aromatic group proton appears. The pyran ring (CH) attached to the ring has a signal at ppm (6.84) attributed to the protonation of the ( $\text{NH}_2$ ) group, and the signal appears at. The signal appears at the ppm site (3.40) attributed to the proton of the group (2CH) attached to the aromatic ring (3.40) and at the ppm site (2.51) attributed to the solvent (DMSO- $d_6$ ), as shown in Figure 4




Figure (4): The  $^1\text{H}$ -NMR spectrum of the compound [MH36]

The signal for compound [MH36] may be seen in its  $^{13}\text{C}$ -NMR spectrum at a ppm point (147.35), which is located distant from the group carbon ( $=\text{CO}-\text{O}$ ) of the compound. The signal that occurs at the ppm location (159.61) is assigned to the group carbon ( $=\text{C}-\text{NH}_2$ ), the signal that appears at the ppm position (122.76-157.86) is credited to the two ring



1H36

Chemical shifts (ppm): 159.61, 157.86, 147.35, 143.21, 136.64, 135.73, 131.48, 128.29, 127.78, 112.58, 39.78, 39.75, 39.72, 39.69, 39.66, 39.63, 39.60, 39.57, 39.54, 39.51, 39.48, 39.45, 39.42, 39.39, 39.36, 39.33, 39.30, 39.27, 39.24, 39.21, 39.18, 39.15, 39.12, 39.09, 39.06, 39.03, 39.00, 38.97, 38.94.

Parameter	Value
1 Data File Name	mohammed
2 Title	MH36
3 Comment	M.m (MH36) 13CNMR in DMSO-d6 at 298K 02.02.03.
4 Origin	UXNMR, Bruker Analytische Messtechnik GmbH
5 Owner	root
6 Site	
7 Instrument	spect
8 Author	
9 Solvent	DMSO
10 Temperature	671.0
11 Pulse Sequence	zgpg30
12 Experiment	ID
13 Probe	5 mm QNP 1H/ 15N/ 13C/ 31P ZGPG02/03058
14 Number of Scans	512
15 Receiver Gain	4597.6
16 Relaxation Delay	1.0000
17 Pulse Width	12.1250
18 Presaturation Frequency	
19 Acquisition Time	1.0912
20 Acquisition Date	2023-05-23T15:24:00
21 Modification Date	2023-05-24T18:25:18
22 Class	
23 Spectrometer Frequency	125.77
24 Spectral Width	30030.0
25 Lowest Frequency	-2441.2
26 Nucleus	13C
27 Acquired Size	32768
28 Spectral Size	65536
29 Digital Resolution	0.46

### 3.3 Evaluation of the Biological Activity of Prepared Compounds

These bacteria were chosen for their medical importance, as they cause many diseases and have different resistance to antibiotics. The bioavailability of many of the prepared compounds was evaluated using etching methods and antibiotic-level measurements [17,18]. The results showed that different ratios of Gram-positive and Gram-negative compounds could inhibit the growth of bacteria, as shown in Table 3 [19,20].

## 4. Conclusion

## REFERENCES

- Central Asian Journal of Medical and Natural Science 2024, 5(4), 130-138. <https://cajmns.centralasianstudies.org/index.php/CAJMNS>

- [2] N. Kerru, L. Gummidi, S. Maddila, and S. B. Jonnalagadda, "A review of recent advances in the green synthesis ofazole-and pyran-based fused heterocycles using MCRs and sustainable catalysts," *\*Current Organic Chemistry\**, vol. 25, no. 1, pp. 4-39, 2021.
- [3] Z. Tashrifi, M. Mohammadi-Khanaposhtani, H. Hamedifar, B. Larijani, S. Ansari, and M. Mahdavi, "Synthesis and pharmacological properties of polysubstituted 2-amino-4H-pyran-3-carbonitrile derivatives," *\*Molecular Diversity\**, vol. 24, no. 4, pp. 1385-1431, 2020.
- [4] A. S. Waghmare, S. S. Pandit, and D. M. Suryawanshi, "DABCO catalyzed green and efficient synthesis of 2-Amino-4H-Pyrans and their biological evaluation as antimicrobial and anticancer agents," *\*Combinatorial Chemistry & High Throughput Screening\**, vol. 21, no. 4, pp. 254-261, 2018.
- [5] M. Maddahi, S. Asghari, and G. F. Pasha, "A facile one-pot green synthesis of novel 2-amino-4 H-chromenes: antibacterial and antioxidant evaluation," *\*Research on Chemical Intermediates\**, vol. 49, no. 1, pp. 253-272, 2023.
- [6] S. B. Lagu et al., "Antitubercular activity assessment of fluorinated chalcones, 2-aminopyridine-3-carbonitrile and 2-amino-4H-pyran-3-carbonitrile derivatives: In vitro, molecular docking and in-silico drug likeliness studies," *\*PloS one\**, vol. 17, no. 6, pp. e0265068, 2022.
- [7] S. D. Hadiyal, N. D. Parmar, P. L. Kalavadiya, J. N. Lalpara, and H. S. Joshi, "Microwave-Assisted Three-Component Domino Synthesis of Polysubstituted 4H-Pyran Derivatives and Their Anticancer Activity," *\*Russian Journal of Organic Chemistry\**, vol. 56, no. 4, pp. 671-678, 2020.
- [8] M. Suthar, J. Kumbhani, and K. E. Y. U. R. d BHATT, "Pyran Heterocyclic Compound as the Prosperous Scaffolds for Biological Sensor (A-Review)," *\*Oriental Journal Of Chemistry\**, vol. 37, no. 6, pp. 1280-1286, 2021.
- [9] W. M. Al-Joboury, K. A. Al-Badrany, and N. J. Asli, "N-alkylation of substituted 2-amino benzothiazoles by 1, 4-bis (bromo methyl) benzene on mixed oxides at room temperature and study their biological activity," in *\*AIP Conference Proceedings\**, vol. 2394, no. 1, pp. 040054, Nov. 2022.
- [10] J. N. Saleh and A. Khalid, "Synthesis, characterization and biological activity evaluation of some new pyrimidine derivatives by solid base catalyst AL<sub>2</sub>O<sub>3</sub>-OBa," *\*Central Asian Journal of Medical and Natural Science\**, vol. 4, no. 4, pp. 231-239, 2023.
- [11] M. J. Saleh, J. N. Saleh, and K. Al-Badrany, "Preparation, characterization, and evaluation of the biological activity of pyrazoline derivatives prepared using a solid base catalyst," *\*EUROPEAN JOURNAL OF MODERN MEDICINE AND PRACTICE\**, vol. 4, no. 7, pp. 25-32, 2024.
- [12] A. W. A. S. Talluh, M. J. Saleh, J. N. Saleh, K. Al-Badrany, and H. mohammed saleh Al-Jubori, "Preparation, characterization, and evaluation of the biological activity of new 2, 3-dihydroquinazoline-4-one derivatives," *\*EUROPEAN JOURNAL OF MODERN MEDICINE AND PRACTICE\**, vol. 4, no. 4, pp. 326-332, 2024.
- [13] A. W. A. S. Talluh, "Preparation, Characterization, Evaluation of Biological Activity, and Study of Molecular Docking of Azetidine Derivatives," *\*Central Asian Journal of Medical and Natural Science\**, vol. 5, no. 1, pp. 608-616, 2024.
- [14] A. W. A. S. Talluh, J. N. Saleh, and M. J. Saleh, "Preparation, Characterization and Evaluation of Biological Activity and Study of Molecular Docking of Some New Thiazolidine Derivatives," *\*World of Science: Journal of Modern Research Technologies\**, vol. 3, no. 2, pp. 49-57, 2024.
- [15] K. A. Al-Badrany, "THE USE OF 2-AMINOPYRAZINE AS A BASIC NUCLEOPHILE FOR THE PREPARATION OF NEW DERIVATIVES OF THE 5, 6-DIHYDROPYRIDINE-2 (1H)-YLIDENE) CYANAMIDE RING, THEIR DIAGNOSIS, AND EVALUATION OF THEIR BACTERIAL EFFICACY," *\*EUROPEAN JOURNAL OF MODERN MEDICINE AND PRACTICE\**, vol. 4, no. 5, pp. 508-518, 2024.
- [16] M. J. Saleh and K. A. Al-Badrany, "Preparation, and characterization of new 2-oxo pyran derivatives by AL<sub>2</sub>O<sub>3</sub>-OK solid base catalyst and biological activity evaluation," *\*Central Asian Journal of Medical and Natural Science\**, vol. 4, no. 4, pp. 222-230, 2023.
- [17] S. H. Abdullah, M. M. Salih, and A. Khalid, "Synthesis, Characterization and Antibacterial Evaluation of Novel Thiazolidine Derivatives."



- 
- [18] A. S. T. Abdul Wahed, "Preparation and Evaluation of Bacterial Activity and Study of the Crystalline Properties of Some 1, 3-Oxazepine-4, 7-Dione Derivatives," *\*Central Asian Journal of Theoretical and Applied Sciences\**, vol. 5, no. 2, pp. 15-26, 2024.
- [19] A. H. Dalaf, M. J. Saleh, and J. N. Saleh, "GREEN SYNTHESIS, CHARACTERIZATION, AND MULTIFACETED EVALUATION OF THIAZOLIDINONE DERIVATIVES: A STUDY ON BIOLOGICAL AND LASER EFFICACY," *\*EUROPEAN JOURNAL OF MODERN MEDICINE AND PRACTICE\**, vol. 4, no. 7, pp. 155-168, 2024.
- [20] A. W. A. S. Talluh, M. J. Saleh, and J. N. Saleh, "Preparation, Characterisation and Study of the Molecular Docking of Some Derivatives of the Tetrazole Ring and Evaluation of their Biological Activity," *\*World of Medicine: Journal of Biomedical Sciences\**, vol. 1, no. 7, pp. 15-23, 2024.