



Article

Synthesis, Characterization and Evaluation of Bacterial Efficacy and Study of Molecular Substrates of Cobalt (II) Complex [Co (2-(benzo[d]thiazol-2-yl)oxy) acetohydrazide) (H₂O) (Cl₂)]

Mahmoud Mehdi Saleh¹, Fahad Farhan Rokan², Jamil Nadhem Saleh³, Mohammed Jwher Saleh⁴

1. Tikrit University, College of Education for Pure Science, Department of Chemistry
2. Kirkuk, Ministry of Education, Department of Chemistry
- 3,4. Salah al-Din Education Directorate, Iraq

* Correspondence: jamelnadhemsaleh@gmail.com

Abstract: Due to the importance of medical and biological cobalt complexes, in this research, a new complex of cobalt complexes was prepared [Co (2-(benzo[d]thiazol-2-yl)oxy)acetohydrazide)(H₂O)(Cl₂)] using 2-(benzo[d]thiazol-2-yl)oxyacetohydrazide as a basic alkand. the ligand was prepared from the reaction of the ester with aqueous hydrazine. , using physical techniques (color, melting point, *R_f*) and spectral methods (FT-IR infrared spectrum, carbon-Proton NMR spectrum ¹H-NMR , ¹³C-NMR, SEM & XRD) the correctness of the compositions of the prepared compounds was confirmed , and then the bioeffectiveness of the series of compounds was studied on two types of Gram-positive and Gram-negative bacteria (*Staphylococcus aureus* and *Escherichia coli*), And the study of the molecular Docking of the complex and the ligand on *Pseudomonas aeruginosa* bacteria and the calculation of the binding energy and RMSD values.

Keywords: Hydrazide, Benzothiazole, Complexes, Molecular Docking, Biological Activity

Citation: Mahmoud Mehdi Saleh. Synthesis, Characterization and Evaluation of Bacterial Efficacy and Study of Molecular Substrates of Cobalt (II) complex [Co (2-(benzo[d]thiazol-2-yl)oxy) acetohydrazide) (H₂O) (Cl₂)]. Central Asian Journal of Medical and Natural Science 2024, 5(4), 198 – 211.

Received: 10th Jul 2024
Revised: 11th Aug 2024
Accepted: 24th Sep 2024
Published: 27th Oct 2024



Copyright: © 2024 by the authors. Submitted for open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>)

1. Introduction

Recently, fused heterocyclic compounds have piqued the interest of medicinal chemists due to their considerable contribution to the biological profile of medications, such as anticancer treatments [1, 2]. Researchers have paid close attention to benzothiazole alternatives because of their widespread usage in pharmaceutical and agricultural chemistry. As a result, the research of benzothiazole derivatives is very significant right now because of their crucial biological and biophysical features, such as antimicrobials [3, 4] and antifungal agents [5]. Several substituted benzothiazole compounds have been found as effective antihelmintic agents [6]. Amino benzothiazole compounds can exhibit a wide range of biological actions, including anti-Parkinson's (a nervous system disorder), dopamine antagonists, antimicrobial, and antihistamine properties [7].

Benzothiazole hydrazine is a bicyclic structure that belongs to the hydrazide group. Because of their multi-reaction chemical composition, benzothiazole derivatives have been extensively researched and are regarded as the nucleus structure for the synthesis of novel benzothiazole derivatives [8]. They have demonstrated a wide range of chemotherapeutic activities, including antimicrobials [9], anti-pain [10], antitumor agents [11], anti-inflammatory [12], anti-cancer [13], antibacterial [14], antifungal [15], anti-HIV [16], is a

potent animal growth stimulator [17]. Mineral complexes, in addition to a variety of benzothiazole derivatives, have antibacterial and antifungal properties [18]. Many drugs' activities have been shown to be increased when the chemical is regarded a ligand in combination with minerals [19].

Hydrazides are versatile compounds that can be used to create heterocyclic Xsystems. Furthermore, because of their diverse biological properties, such as antidiabetic [20], antimalarial [21], anticancer [22], antimicrobial [23], antioxidant [24], anti-inflammatory [25], and anxiolytic [26], hydrazide-hydrazones continue to be of great interest in the field of medicinal chemistry.

2. Materials and Methods

Material

Without any additional purification, all of the compounds utilized in this research investigation were acquired from BDH, Fluka, and Aldrich.

Devices Used:

Melting points were measured using a Thermoelectric Melter 9300 at a scale of 400-4000 cm⁻¹, a Shimadzu FT-IR 8400S spectrophotometer, and a Bruker apparatus running at 400 MHz for ¹H-NMR and ¹³C-NMR spectra, respectively. Fluka silica gel plates were used in thin-layer chromatography (TLC) and had a thickness of 0.2 mm. UV light was used to restore visibility after the plates had been activated with fluorescent silica gel. The scanning electron microscope (SEM) (Czech Republic/Belsorp Mini II/TE SCAN) was utilized.

Preparation of 2-(benzo[d]thiazol-2-yloxy)acetohydrazide (MH1) [27]

A solution of ethyl 2-(benzo[d]thiazol-2-yloxy) acetate (0.01 mol) and hydrazine hydrate 99% (0.2 mol) in abs ethanol (50 mL) was refluxed for three hours. Excess ethanol was eliminated through distillation. Upon cooling, 2-(benzo[d]thiazol-2-yloxy) acetohydrazide began to separate. It was filtered and recrystallized with ethanol, yielding an 84% product with a melting point of 190 degrees Celsius and a faint yellow color.

Preparation of Complex (MH2) [28]


00.2 g dissolved from 2-(benzo[d]thiazol-2-yloxy)acetohydrazide in 10mL of absolute ethanol was mixed with 0.210 g of CoCl₂. 6H₂O dissolved in 10mL of distilled water and heated the mixture with a release at a temperature of 150C for three hours, then the solution was cooled, concentrated and recrystallized in the resulting had a pink color and a melting point of 300

Biological activity study:

Biological activity has been measured using the Kirby Bauer movement's propagation method, which entails spreading 0.1 ml of bacterial solution on Nutrient Broth plates and allowing them to absorb the fluid for five minutes [29, 30]. To make holes in each dish, a five mm cork and a cork plunger were utilized [31,32]. After that, 0.1 ml of the fourth hole's generated solutions-which used ciprofloxacin as a reference sample-were incubated for twenty-four hours at 37 °C. The inhibitory zone widths around each hole have been measured in millimeters using Prescott's method [33,34,35].

3. Results and Discussion

This study involved the preparation of several compounds, as the scheme(1) illustrates.



Scheme 1. Path of the Ready Combination (MH1, MH2)

Characterization of hydrazide [36]

FT-IR(KBr): (3224) ν (N-H), (2970) ν (C-H), (3083) ν (Ar-H), (1677) ν (C=O), (1604) ν (C=N), (3309,3390) ν (NH₂), (1475,1550) ν (C=C), as shown in the figure (1). ¹H-NMR (DMSO-d₆) δ (PPm): (4.37) (s, NH₂), (9.47) (s, NH), (4.85) (s, CH₂), (7.18-7.68) (m, Ar-CH). as shown in figure (2). ¹³C-NMR (DMSO-d₆) δ (PPm) : (43.76) (CH₂), (165.84) (C=O), (169.56) (C=N), (111.85-137.75) (m, Ar-CH). as shown in figure (3).




Figure. The Infrared Spectrum of the Compound (MH1)




Figure (4): The infrared spectrum of the compound (MH2)




Figure (5): The ¹H-NMR spectrum of the compound (MH2)




Figure (6): The ¹³C-NMR spectrum of the compound (MH2)

The Results of SEM:

Scanning electron microscope (SEM) technology is based on the use of a high-energy focused electron beam, which in turn interacts with the surface of the sample exposed to it, resulting in obtaining topographic images, as well as generating a different set of signals on the surface of the solid sample under test, and this technique can provide a lot of information about the sample surface, and this technique depends mainly on the area of the spot that is exposed to the electronic beam as well as the extent of its impact on it, where dot scanning is generally used to scan the electron beam and then the signal is combined with its location to produce a specific image of the surface of the sample under test [38,39]

Scanning electron microscopy imaging (Complex 2) showed nanoparticles of irregular shapes and sizes, ranging in diameters from 33-334 Nm, and also showed many interstitial distances between minutes, indicating high porosity and high absorbability. As shown figure (7).




Figure 7. Scanning Electron Imaging of the MH2 Complex

The Results of XRD [40]:

The complex (MH2) has shown beams of high and medium intensity in the diffraction spectrum within the range of 50-10 O, which indicates the presence of a crystal structure of the compound, as well as wide beams, i.e. high FWHM values, indicate the small nanomicros and high porosity of the complex, which was calculated by the Scherrer equation above and the minute volume rate was equal to (11.44 nm), see Figure (8), Table (1) shows the data of the diffraction spectra of the measured complexes as well as the.




Figure 8. Diffraction Spectra of the Complex (MH2)

Table 1. Data of Doffraction Spectra of the Complex (MH2) as well as the Rate of Minute Volumes According to the Scherrer Equation

Compounds	2 θ	FWHM	Intensity	d spacing	D (nm)	Avg. D (nm)
[MM2)]	10.5886	0.3936	180.64	8.35508	21.17653	19.2916
	13.457	1.1808	57.5	6.57993	7.077475	
	18.8696	0.246	269.92	4.70299	34.20054	
	23.6523	0.3444	309.27	3.76172	24.62111	
	25.874	0.3936	141.28	3.44354	21.63536	
	29.3689	0.2952	194.18	3.04122	29.06425	
	41.4743	0.7872	77.29	2.17729	11.27345	
	44.221	1.1808	50.61	2.04822	7.586654	
	48.4092	0.492	109.67	1.88035	18.49489	
	75.5581	0.5904	61.21	1.25843	17.78572	

Evaluation of the Biological Activity:

Certain compounds (MH1, MH2) were evaluated against several bacterial strains using the cup plate agar diffusion method, including gram-negative *Escherichia coli* and gram-positive *Staphylococcus aureus* [41, 42, 43]. After eight hours of incubation at 37 °C, the microbial cultures were diluted with 0.8% sterile saline [44,45,46]. The medicines were dissolved in DMSO with a concentration of 100 g/mL. Amoxicillin was employed as a negative control in this experiment. The diameter of the ring encircling the active disk in bacterial growth inhibition was utilized to determine biological activity, as shown in Table (2)[47, 48, 49].

Table 2. Inhibitory Effectiveness of Some Prepared Compound (MH1, MH2) and Control Treatments (Antibiotics) on the Growth of a Number of Positive and Negative Bacteria

Comp. No.	<i>Escherichia coli</i>			<i>Staphylococcus aureus</i>		
	0.001	0.01	0.1	0.001	0.01	0.1
MH1	23	14	12	30	45	50
MH2	19	26	28	10	25	30
Amoxicillin	10	16	24	10	20	20




Figure 9. Biological Effect of Some Bacterial of *Staphylococcus Aureus* and Inhibition Zone for three Concentration




Figure 10. Biological Effect of Some Bacterial of Escherichia Coli and Inhibition Zone for Three Concentration

Results of Molecular Docking Study for MH2 [50]:

The molecular docking of the complex (MH2) was studied on a single line it is a bacterium *Pseudomonas aeruginosa*, using the MOE program (2014), where the energy reduction process was completed for vehicles under in order to get the most stable vacuum body (the least energy hindrance), then download Synthesis of *Pseudomonas aeruginosa* bacteria from the World Bank protein site (receptor X3R6), a calculator was used Personality. The binding energy of (MH1) was -4.69, with an RMSD value of 3.13. The binding energy of the complex (MH2) was -6.02, with an RMSD value of 2.08.

The investigation of the molecular adhesion of the revealed the number and types of bonds that these produced derivatives create with amino acid residues in the active site, generating a number of connections.

The study showed that the compound (MH1) interacts with amino acid residues that are present in the active site by forming two types of bonds, hydrogen bonds that bind amino acid residues. 406Tyr is at the active site with the electron pair of the oxygen atom, the carbonyl group with a length of 2.82, and hydrogen bonds that bind the residues of amino acids 331. The Arg is placed in the active site with the electron pair of a sulfur atom with a length of 3.33 and its PI-type interaction with a benzene ring, a number of amino acids impacted by vandervalls forces, and binding energy values of -4.69 were also determined. Figure (11) depicts the findings of molecular anchoring and the interaction of the synthesized chemical MH1 with the receptor X3R6 of a single line of *Pseudomonas aeruginosa* bacteria.





Figure 11. Shows a two-dimensional and Three-Dimensional Representation of the Results of Molecular Docking and the Association Between the Compound (MH1) Prepared Receptor X3R6 Single Line of *Pseudomonas Aeruginosa* Bacteria

The researchers observed that the chemical (MH2) interacts with amino acid residues in the active site by creating two types of alkyl-Pi ligands that bind to them. 487thr, which is at the active site with the aromatic ring's electron pairs and a number of amino acids affected by vandervalz forces, was discovered to have a binding energy of -6.02. Figure 12 depicts a two-dimensional and three-dimensional representation of the molecular docking data, as well as the interaction between the generated chemical (MH2) and the receptor 6X3R one line of the *Pseudomonas aeruginosa* bacteria.






Figure 12. Shows Two-Dimensional and Three-Dimensional Representation of the Results of Molecular Docking and the Association Between the Compound (MH1) Prepared Receptor X3R6 Single Line of *Pseudomonas Aeruginosa* Bacteria

4. Conclusion

The initial confirmation of the prepared compounds was made through color and temperature changes, and the final confirmation was made through the appearance of distinct bands in the FT-IR spectrum, as well as the appearance of the expected signals for the prepared compounds in the C13 H1 NMR spectrum. In addition, the compounds demonstrated strong inhibitory efficacy against bacteria as compared to a control sample.

REFERENCES

- [1] Abbas, E. M., Amin, K. M., El-Hamouly, W. S., Dawood, D. H., & Abdalla, M. M. (2015). Synthesis, anti-inflammatory and antinociceptive activity of some novel benzothiazole derivatives. *Research on chemical Intermediates*, 41, 2537-2555.
- [2] Abdul Wahed, A. S. T. (2024). Preparation and Evaluation of Bacterial Activity and Study of the Crystalline Properties of Some 1, 3-Oxazepine-4, 7-Dione Derivatives. *Central Asian Journal of Theoretical and Applied Sciences*, 5(2), 15-26.
- [3] Aftan, M. M., Salih, H. K., & Talloh, A. A. (2021). Synthesis of new mesogenic Schiff bases ether with polar halogen Substituent and study their liquid crystalline properties. *Journal of Education and Scientific Studies*, 5(17).
- [4] Al Rashidy, A. A. M., Al Badrany, K. A., & Al Garagoly, G. M. (2020, August). Spectrophotometric determination of sulphamethoxazole drug by new pyrazoline derived from 2, 4-dinitro phenyl hydrazine. In *Materials Science Forum* (Vol. 1002, pp. 350-359). Trans Tech Publications Ltd.
- [5] AL SOUD, Y. A., AL SADONI, H. H., Saeed, B., Jaber, I. H., MO, B. K., AL MASOUDI, N. A., ... & Loddo, R. (2008). Synthesis and in vitro antiproliferative activity of new benzothiazole derivatives. *Arkivoc*, 2008(15), 225-238.
- [6] Alasadi, Y. K., Jumaa, F. H., Dalaf, A. H., Shawkat, S. M., & Mukhlif, M. G. (2022). Synthesis, Characterization, and Molecular Docking of New Tetrazole Derivatives as Promising Anticancer Agents. *Journal of Pharmaceutical Negative Results*, 13(3), 513-522.
- [7] Alazzawi, S. S., & Alasadi, Y. K. (2023). Synthesis and study of the biological activity and molecular Docking of 2-hydrazinobenzothiazole derivatives. *Multi-Knowledge Electronic Comprehensive Journal For Education & Science Publications (MECSJ)*, (62).

- [8] Al-Badrany, A. S. S., & Al, R. J. S. A. D. (2023). The Impact of the Digital Economy on International Trade, the Case of Egypt for the Period (1990-2020). *Integrated Journal for Research in Arts and Humanities*, 3(2), 163-173.
- [9] Al-Badrany, K. A. (2024). THE USE OF 2-AMINOPYRAZINE AS A BASIC NUCLEOPHILE FOR THE PREPARATION OF NEW DERIVATIVES OF THE 5, 6-DIHYDROPYRIDINE-2 (1H)-YLIDENE) CYANAMIDE RING, THEIR DIAGNOSIS, AND EVALUATION OF THEIR BACTERIAL EFFICACY. *EUROPEAN JOURNAL OF MODERN MEDICINE AND PRACTICE*, 4(5), 508-518.
- [10] Al-Badrany, K. A., Mohammed, A. S., & Alasadi, Y. K. (2019). Synthesis of some new 1, 3, 4-oxadiazole compounds derived from 1H-imidazole and study their biological activity. *Eurasian Journal of Biosciences*, 13(1), 501-507.
- [11] Al-Hadidi, O. A. F., Al-Badrany, K. A., & Al-Bajari, S. A. (2022). Synthesis some of thiazepine compounds from 2-carboxyaldehyde-5-methyl thiophene and study their biological activity on infected male rats epileptic. *Journal of Education and Scientific Studies*, 2(20).
- [12] Al-Joboury, W. M., Al-Badrany, K. A., & Asli, N. J. (2022, November). N-alkylation of substituted 2-amino benzothiazoles by 1, 4-bis (bromo methyl) benzene on mixed oxides at room temperature and study their biological activity. In *AIP Conference Proceedings* (Vol. 2394, No. 1). AIP Publishing.
- [13] Al-Tufah, M. M., Jasim, S. S., & Al-Badrany, K. A. (2020). Synthesis and Antibacterial Evaluation of some New Pyrazole Derivatives. *Prof. (Dr) RK Sharma*, 20(3), 178.
- [14] Apelt, J., Grassmann, S., Ligneau, X., Pertz, H. H., Ganellin, C. R., Arrang, J. M., ... & Stark, H. (2005). Search for histamine H3 receptor antagonists with combined inhibitory potency at N τ -methyltransferase: ether derivatives. *Die Pharmazie-An International Journal of Pharmaceutical Sciences*, 60(2), 97-106.
- [15] Catalano, A., Carocci, A., Defrenza, I., Muraglia, M., Carrieri, A., Van Bambeke, F., ... & Franchini, C. (2013). 2-Aminobenzothiazole derivatives: search for new antifungal agents. *European Journal of Medicinal Chemistry*, 64, 357-364.
- [16] Dalaf, A. H., Saleh, M. J., & Saleh, J. N. (2024). GREEN SYNTHESIS, CHARACTERIZATION, AND MULTIFACETED EVALUATION OF THIAZOLIDINONE DERIVATIVES: A STUDY ON BIOLOGICAL AND LASER EFFICACY. *EUROPEAN JOURNAL OF MODERN MEDICINE AND PRACTICE*, 4(7), 155-168.
- [17] Dash, D. C., Mahapatra, A., Mohapatra, R. K., Ghosh, S., & Naik, P. (2008). Synthesis and characterization of dioxouranium (VI), thorium (IV), oxozirconium (IV) and oxovanadium (IV) complexes with 1, 11-dihydroxy-1, 4, 5, 7, 8, 11-hexaaza-2, 3, 9, 10-tetramethyl-1, 3, 8, 10-decatetraene-6-thione and their derivatives with chloroacetic acid.
- [18] Duong, T. H., Devi, A. P., Huynh, N. V., Sichaem, J., Tran, H. D., Alam, M., ... & Nguyen, T. C. (2020). Synthesis, α -glucosidase inhibition, and molecular docking studies of novel N-substituted hydrazide derivatives of atranorin as antidiabetic agents. *Bioorganic & Medicinal Chemistry Letters*, 30(17), 127359.
- [19] Gurram, S. R., & Azam, M. A. (2021). Design, synthesis, antibacterial evaluation and molecular docking studies of some newer benzothiazole containing aryl and alkaryl hydrazides. *Chemistry & Biodiversity*, 18(7), e2100117.
- [20] Holobova, E., & Peijessy, A. (1986). synthesis and properties of 2-hydrazinobenzothiazole derivatives. *Chem. Papers*, 40(6), 791-796.
- [21] Huang, G., Cierpicki, T., & Grembecka, J. (2023). 2-Aminobenzothiazoles in anticancer drug design and discovery. *Bioorganic chemistry*, 135, 106477.
- [22] Huang, S. T., Hsei, I. J., & Chen, C. (2006). Synthesis and anticancer evaluation of bis (benzimidazoles), bis (benzoxazoles), and benzothiazoles. *Bioorganic & medicinal chemistry*, 14(17), 6106-6119.
- [23] Irzogi, A. A., Salih, M. M., Jirjes, H. M., & Mensoor, M. K. (2020). Synthesis, Characterization, and Antibacterial Activity of Complexes of Hg (II) with Mixtures of 3-Hydrazonoindolin-2-one and Diphosphine, or Diimine Ligands. *Russian Journal of General Chemistry*, 90, 1069-1073.
- [24] Jabeen, M., Ahmad, S., Shahid, K., Sadiq, A., & Rashid, U. (2018). Ursolic acid hydrazide based organometallic complexes: synthesis, characterization, antibacterial, antioxidant, and docking studies. *Frontiers in Chemistry*, 6, 55.
- [25] Jirjes, H. M., Irzogi, A. A., Al-Doori, L. A., Alheety, M. A., & Singh, P. K. (2022). Nano cadmium (II)-benzyl benzothiazol-2-ylcarbomodithioate complexes: Synthesis, characterization, anti-cancer and antibacterial studies. *Inorganic Chemistry Communications*, 135, 109110.
- [26] Khalil, M. I., & Khalal, Q. Z. (2021, March). Synthesis and characterization of new compounds derived from 2-hydrazinobenzothiazole and evaluated their antibacterial activity. In *Journal of Physics: Conference Series* (Vol.

- 1853, No. 1, p. 012007). IOP Publishing.
- [27] Luo, B., Li, D., Zhang, A. L., & Gao, J. M. (2018). Synthesis, antifungal activities and molecular docking studies of benzoxazole and benzothiazole derivatives. *Molecules*, 23(10), 2457.
- [28] Mohamed, S. A., Hussein, M. S., & Al-badrany, K. A. (2022). Synthesis and characterization of pyrazolines and oxazapine derivatives using chalcones as precursor and evaluation of their biological activity. *Samarra Journal of Pure and Applied Science*, 4(4).
- [29] Morsy, M. A., Ali, E. M., Kandeel, M., Venugopala, K. N., Nair, A. B., Greish, K., & El-Daly, M. (2020). Screening and molecular docking of novel benzothiazole derivatives as potential antimicrobial agents. *Antibiotics*, 9(5), 221.
- [30] Mutlag, Z. J., & Irzoqi, A. A. (2024). SYNTHESIS, DIAGNOSIS, THERMAL AND SURFACE STUDY AND EVALUATION OF BACTERIAL EFFICACY OF SOME CADMIUM (II) COMPLEXES WITH A COMBINATION OF 4-METHOXYBENZALENE (HYDRAZINEYL AND PHOSPHINES) LIGANDS. *EUROPEAN JOURNAL OF MODERN MEDICINE AND PRACTICE*, 4(6), 544-555.
- [31] Nagarajan, S. R., De Crescenzo, G. A., Getman, D. P., Lu, H. F., Sikorski, J. A., Walker, J. L., ... & Blystone, L. (2003). Discovery of novel benzothiazolesulfonamides as potent inhibitors of HIV-1 protease. *Bioorganic & medicinal chemistry*, 11(22), 4769-4777.
- [32] Pan, X., Chen, L., Xu, W., Bao, S., Wang, J., Cui, X., ... & Chen, R. (2021). Activation of monoaminergic system contributes to the antidepressant-and anxiolytic-like effects of J147. *Behavioural Brain Research*, 411, 113374.
- [33] Pathak, N., Rath, E., Kumar, N., Kini, S. G., & Rao, C. M. (2020). A review on anticancer potentials of benzothiazole derivatives. *Mini reviews in medicinal chemistry*, 20(1), 12-23.
- [34] Popiołek, Ł. (2021). Updated Information on Antimicrobial Activity of Hydrazone-Hydrazones. *International Journal of Molecular Sciences*, 22(17), 9389.
- [35] Ramírez, H., Fernandez, E., Rodrigues, J., Mayora, S., Martínez, G., Celis, C., ... & Charris, J. (2021). Synthesis and antimalarial and anticancer evaluation of 7-chloroquinoline-4-thiazoleacetic derivatives containing aryl hydrazone moieties. *Archiv der Pharmazie*, 354(7), 2100002.
- [36] Reddy, A. K., & Kathale, N. E. (2017). Synthesis and anti-inflammatory activity of hydrazones bearing biphenyl moiety and vanillin based hybrids. *Oriental Journal of Chemistry*, 33(2), 971.
- [37] Sadeek, G. T., Saeed, Z. F., & Saleh, M. Y. (2023). Synthesis and Pharmacological Profile of Hydrazone Compounds. *Research Journal of Pharmacy and Technology*, 16(2), 975-982.
- [38] Saleh, J. N., & Khalid, A. (2023). Synthesis, characterization and biological activity evaluation of some new pyrimidine derivatives by solid base catalyst Al_2O_3 -OBa. *Central Asian Journal of Medical and Natural Science*, 4(4), 231-239.
- [39] Saleh, M. J., & Al-Badrany, K. A. (2023). Preparation, characterization of new 2-oxo pyran derivatives by Al_2O_3 -OK solid base catalyst and biological activity evaluation. *Central Asian Journal of Medical and Natural Science*, 4(4), 222-230.
- [40] Saleh, M. J., Saleh, J. N., & Al-Badrany, K. (2024). PREPARATION, CHARACTERIZATION, AND EVALUATION OF THE BIOLOGICAL ACTIVITY OF PYRAZOLINE DERIVATIVES PREPARED USING A SOLID BASE CATALYST. *EUROPEAN JOURNAL OF MODERN MEDICINE AND PRACTICE*, 4(7), 25-32.
- [41] Saleh, R. H., Rashid, W. M., Dalaf, A. H., Al-Badrany, K. A., & Mohammed, O. A. (2020). Synthesis of some new thiazolidinone compounds derived from schiff bases compounds and evaluation of their laser and biological efficacy. *Ann Trop & Public Health*, 23(7), 1012-1031.
- [42] Shcherbakov, V. V., Artemkina, Y. M., Akimova, I. A., & Artemkina, I. M. (2021). Dielectric characteristics, electrical conductivity and solvation of ions in electrolyte solutions. *Materials*, 14(19), 5617.
- [43] Singh, M. K., Tilak, R., Nath, G., Awasthi, S. K., & Agarwal, A. (2013). Design, synthesis and antimicrobial activity of novel benzothiazole analogs. *European journal of medicinal chemistry*, 63, 635-644.
- [44] Suresh, C. H., Rao, J. V., Jayaveera, K. N., & Subudhi, S. K. (2011). Synthesis and anthelmintic activity of 3 (2-hydrazino benzothiazoles)-substituted indole-2-one. *IJPR*, 2, 257-61.
- [45] Talluh, A. W. A. S. (2024). Preparation, Characterization, Evaluation of Biological Activity, and Study of Molecular Docking of Azetidine Derivatives. *Central Asian Journal of Medical and Natural Science*, 5(1), 608-616.
- [46] Talluh, A. W. A. S., Saleh, J. N., & Saleh, M. J. (2024). Preparation, Characterization and Evaluation of Biological Activity and Study of Molecular Docking of Some New Thiazolidine Derivatives.

- [47] Talluh, A. W. A. S., Saleh, M. J., & Saleh, J. N. (2024). Preparation, Characterisation and Study of the Molecular Docking of Some Derivatives of the Tetrazole Ring and Evaluation of their Biological Activity. *World of Medicine: Journal of Biomedical Sciences*, 1(7), 15-23.
- [48] Talluh, A. W. A. S., Saleh, M. J., Saleh, J. N., Al-Badrany, K., & mohammed saleh Al-Jubori, H. (2024). Preparation, characterization, and evaluation of the biological activity of new 2, 3-dihydroquinazoline-4-one derivatives. *EUROPEAN JOURNAL OF MODERN MEDICINE AND PRACTICE*, 4(4), 326-332.
- [49] Umapathi, A., Navya, P. N., Madhyastha, H., Singh, M., Madhyastha, R., Maruyama, M., & Daima, H. K. (2020). Curcumin and isonicotinic acid hydrazide functionalized gold nanoparticles for selective anticancer action. *Colloids and Surfaces A: Physicochemical and Engineering Aspects*, 607, 125484.
- [50] Zaki, A. N. M., Husain, S. S., & Obaid, B. H. (2016). Effect of some derivatives quinoline prepared on some pathogenic bacteria species compared with the 2-hydrazinobenzothiazole. *AGERC*, 805.