



Article

The Association between Pregnant Women's Satisfaction and Antenatal Care Service in Maysan Public Health Centers

Haifaa Kadhim AL-Zaidy¹, Sarab Nasr Fadhil²

¹Community Health Nursing Specialist, MSN, Ministry of Health, Maysan, Iraq, Email: Haifaa.gasem2206m@conursing.uobaghdad.edu.iq

²Department of Maternal and Neonate Nursing, College of Nursing, University of Baghdad, Iraq, Email: sarab@conursing.uobaghdad.edu.iq

* Correspondence: sarab@conursing.uobaghdad.edu.iq

Abstract: Pregnancy is a very important and special event from both social and medical points of view. Maternal satisfaction with antenatal care (ANC) services has been linked to the quality of the provided services and the extent to which specific maternal needs are met. This study aimed to evaluate pregnant women's satisfaction with antenatal care services at primary health care centers in Maysan city. This study aimed to evaluate pregnant women's satisfaction with antenatal care services at primary health care centers in Maysan city. This quantitative descriptive study was conducted from January 2d to February 19th, 2024. The sample of study included a purposeful (nonprobable) sample method and included 390 pregnant women who visited primary health care centers. The study instrument includes participants' sociodemographic characteristics, reproductive characteristics, The Measure of patient satisfaction with the provided care, Data were collected through self-report and analyzed using the statistical package for social science, IBM version 27. The results show that the multiple regression model negatively predicts women's satisfaction with antenatal care services at primary health care centers (p value = .006). This study concludes that antenatal care services at primary health care centers negatively predict women's satisfaction with these services.

Keywords: Association, Antenatal care, Pregnant women, Satisfaction.

Citation: Haifaa Kadhim AL-Zaidy, Sarab Nasr Fadhil. The Association between Pregnant Women's Satisfaction and Antenatal Care Service in Maysan Public Health Centers Central Asian Journal of Medical and Natural Science 2024, 5(4), 399-409.

Received: 10th May 2024

Revised: 11th Jun 2024

Accepted: 24th Jun 2024

Published: 27th Jul 2024



Copyright: © 2024 by the authors. Submitted for open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>)

1. Introduction

Antenatal care (ANC) plays a critical role in preventing maternal morbidity and mortality. It offers a range of services that include administering preventive measures such as immunizations, educating pregnant women on healthy behaviors during pregnancy and identifying and managing potential complications [1].

Moreover, antenatal care (ANC) is an essential health treatment to help mothers and babies live longer lives. The World Health Organization (WHO) defines adequate

prenatal care as at least four health care visits throughout pregnancy for pregnant women who do not experience perinatal difficulties [2].

Prenatal care is also an efficient prenatal care service for reducing complications during pregnancy, increasing birth weights, and decreasing perinatal, neonatal, infant and child mortality [3]. Despite its benefits, the perceptions and experiences of pregnant women significantly influence the utilization of ANC services [4].

Patient satisfaction is defined as the individual's positive evaluation of a distinct dimension of health care. Client satisfaction measures the ability of utilities to meet consumers' suggestions and is a significant factor in the selection of health facilities and the future use of services, which are determined by the individual's perceptions, attitudes, and comparison processes [5].

Satisfaction has been said to be a major predictor of the use of services, as it is essential if clients are to utilize services, comply with treatments and maintain a continuing relationship with practitioners [6]. Women's satisfaction is ultimately used as a gauge of antenatal care quality, which is a crucial aspect of service evaluation [7].

Low satisfaction rates with the quality of care offered at ANC facilities can discourage women from attending ANC visits, leading to missed opportunities for preventive measures and timely management of high-risk pregnancies [8]. Improving women's health and maintaining healthy behaviors before or during pregnancy is particularly important until nearly half of pregnancies (49%) are unintended and correlated with insufficient information about the risk factors that place them at constant risk [9].

The quality of primary health care services is one of the most important areas in the health services sector. There are several people who can be of high value for public health and private health, and these indicators are the most prominent [10]. Iraq is a country with a high maternal mortality and neonatal mortality ratio (35/100 000) and a high neonatal mortality rate (23/1000). Poverty and adverse socioeconomic conditions lead to early marriage and disruption of education, teenage pregnancy and childbirth, high maternal and neonatal mortality with morbidity and more children born to live in these adverse conditions [11].

A significant number of women die globally because of birth and pregnancy-related complications [12]. In addition, pregnancy health care is the key entry point at which a pregnant woman receives a broad range of health promotion and preventive services that support the health of the mother and the baby [13]. Aims and objectives of this study aimed to evaluate pregnant women's satisfaction with antenatal care services at primary health care centers in Maysan city.

2. Materials and Methods

The study Design

A quantitative design (descriptive study). This study was conducted from January 2ed to February 19th, 2024, to assess pregnant women's satisfaction with antenatal care services at primary health care centers in Maysan city.

Ethical considerations.

This research was confirmed by the Committee of Scientific Research at the College of Nursing, University of Baghdad. The official reference number is (2), dated November 22th, 2023. After obtaining the approval of the Ministry of Planning (Central Statistical Organization), official approval was obtained from the Maysan Health Department. Approval was obtained from the targeted (Human Developmental and Training Center)

to obtain official permission to carry out the study at all primary health care centers in Maysan city, Iraq.

Setting of the Study

This study was conducted in primary health care centers in Maysan city.

The Study Samples

The sample of study included a purposeful (nonprobable) sample method and included 390 pregnant women who visited primary health care centers.

The selection criteria were designed as follows:

Inclusion criteria: were all pregnant women who age ≥ 15 years old, primiparous and multiparous who visited primary health care centers.

Exclusion criteria: Non-pregnant woman, Pregnant women who have any mental disability, deaf, those under the age of fifteen years and those who declined participation

Materials & Measures

To evaluate pregnant women's satisfaction with antenatal care, a questionnaire was constructed by the authors. The researcher emailed the authors who constructed the questionnaire. All were contacted and granted permission to use the questionnaire in research. Official permission was given by the corresponding author of the questionnaire [14]. Which mainly consisting of four parts. The demographic data section was designed to obtain the data of participants in the study. These data included 7 items: age, social status, area of residence, educational level, number of pregnancies, PHC visits, and economic status. The data were collected via a self-reported interview questionnaire. Reproductive Information was collected via a 4-item self-reported interview questionnaire: which include current gestational age, number of births, number of abortions, and pregnancy type. Services and management this part consisted of 10 items, including pregnant woman's weight, height, and blood pressure. Moreover, blood tests, ultrasound, urine tests, and physical examination results were also logged. Furthermore, whether the patient was taking an iron or folic acid prescription, or any other medication was also considered. The level of ANC services was scored by rating (1) for the yes answer and (2) for the no answer. Patient satisfaction with the provided care should be assessed to provide a clear image of the quality of the provided care. This part consists of 13 items. The Satisfaction Category for Disagree, Neutral and Agree, Disagree Scored by Rating (1), Neutral Scored by Rating (2) and agree Scored by Rating (3). The total score is 13-39.

The face validity of the study instrument was determined by a panel of 8 experts. They were all faculty members from the College of Nursing/University of Baghdad. To review the content of the questionnaire, those experts were provided with a print of study instrument and asked to review and evaluate the instrument for its content clarity and adequacy.

Samples were collected daily during the official working days. The questionnaire lasted for 10-15 minutes for each pregnant woman. Women were informed that they could interrupt the interview and withdraw whenever they wanted to answer or felt it difficult to complete the questionnaire. Both self-report and interview approaches were used for data collection.

Statistical analysis: Data were analyzed using the statistical package for social science (SPSS) for windows, version 27. The descriptive statistical measure of frequency

and percent were used to describe participants' sociodemographic characteristics. The arithmetic mean and standard deviation were also used.

3. Results

Table 1. Sociodemographic characteristics of Study Sample

Variable	Frequency	Percent
Age (Years): Mean (SD): 24.62 ± 5.66		
15-21	124	31.8
22-28	179	45.9
29-35	70	17.9
36-43	17	4.4
Marital Status		
Married	390	100.0
Residency		
Urban	327	83.8
Rural	63	16.2
The highest level of education attained		
Unable to read and write	51	13.1
Read and write	66	16.9
Elementary	170	43.6
Middle	59	15.1
High	11	2.8
Diploma	8	2.1
Bachelor's degree	25	6.4
Monthly Income (Iraqi Dinar)		
<300.000	121	31.0
301.000 - 600.000	167	42.8
601.000-900.000	52	13.3
901.000-1200.000	50	12.8

SD: Standard deviation

The mean age is 24.62 ± 5.66 ; less than half aged 22-28 years ($n = 179$; 45.9%), followed by those aged 15-21 years ($n = 124$; 31.8%), and those aged 29-35 years ($n = 70$; 17.9%), and those who age 36-43-years ($n = 17$; 4.4%). The study results reveal that all women are married ($N = 390$; 100.0%). The majority reported that they have been living in urban areas ($n = 327$; 83.8%) compared to those who have been living in rural areas ($n = 63$; 16.2%). Concerning the highest least level of education women attained, more than two-fifths are elementary school graduates ($n = 170$; 43.6%), followed by those who read and write ($n = 66$; 16.9%), those who are middle school graduates ($n = 59$; 15.1%), those who are unable to read and write ($n = 51$; 13.1%), those who hold bachelor's degree ($n = 25$; 6.4%), those who are high school graduates ($n = 11$; 2.8%), and those who hold diploma degree ($n = 8$; 2.1%). Regarding family's monthly income, more than two-fifths reported that their monthly income ranges between 301.000 - 600.000 ID ($n = 167$; 42.8%), followed by those whose monthly income is less than 300.000 ID ($n = 121$; 31.0%), those whose monthly income ranges between 601.000-900.000 ID ($n = 52$; 13.3%), and those

whose monthly income ranges between 901.000-1200.000 ID ($n = 50$; 12.8%).

Table 2: Participants' reproductive profile

Variable	Frequency	Percent
Gravidity		
1	171	43.8
2	85	21.8
3-4	82	21.0
5-6	36	9.2
≥ 7	16	4.1
Is this your first antenatal visit at this facility for this pregnancy?		
Yes	183	46.9
No	207	53.1
Current gestational age (Weeks)		
4-13	22	5.6
14-27	259	66.4
28-39	109	28.0
Parity		
0	192	49.2
1	81	20.8
2	56	14.4
3-4	44	11.3
5-6	13	3.3
≥ 7	4	1.0
Type of pregnancy		
Planned	318	81.5
Not planned	72	18.5

The study results display that more than two-fifths have one pregnancy ($n = 171$; 43.8%), followed by those who have two pregnancies ($n = 85$; 21.8%), those who have 3-4 pregnancies ($n = 82$; 21.0%), those who have 5-6 pregnancies ($n = 36$; 9.2%), and those who have seven or more pregnancies ($n = 16$; 4.1%). When questioned about (Is this your first antenatal visit at this facility), more than two-fifths reported that this is not so ($n = 207$; 53.1%) compared to those who reported that this is so ($n = 183$; 46.9%). Concerning current gestational age, this age ranges between 14-27 weeks for most women ($n = 259$; 66.4%), followed by 28-39 weeks ($n = 109$, 28.0%), and 4-13 weeks ($n = 22$; 5.6%). Concerning parity, around half are nulliparous ($n = 192$; 49.2%), followed by those who gave to one birth ($n = 81$; 20.8%), those who gave to two births ($n = 56$; 14.4%), those who gave to 3-4-births ($n = 44$; 11.3%), those who gave birth to 5-6 births ($n = 13$; 3.3%), and those who gave to 7 or more births ($n = 4$; 1.0%). Regarding the type of pregnancy, the majority reported that it is planned ($n = 318$; 81.5%) compared to not planned ($n = 72$; 18.5%).

Table 3: Levels of women's satisfaction with antenatal care services at primary healthcare centers

Satisfaction	Frequency	Percent
Dissatisfied	2	0.5
Somewhat satisfied	331	84.9
Satisfied	57	14.6

The study revealed that most women were somewhat satisfied with antenatal care services at primary healthcare centers (n = 331; 84.9%), followed by those who were satisfied with them (n = 57; 14.6%) and those who were dissatisfied with them (n = 2; 0.5%).

Table 4: Differences in women's satisfaction with antenatal care service at primary health care centers between residency groups

Ranks					Mann-Whitney U	Asymp. Sig. (2-tailed)
	Residency	N	Mean Rank	Sum of Ranks		
Patient Satisfaction	Urban	327	195.09	63793.50	10165.500	.867
	Rural	63	197.64	12451.50		
	Total	390				

The study results reveal that there is no statistically significant difference in women's satisfaction with antenatal care service at primary health care centers between residency groups.

Table 5: Differences in women's satisfaction with antenatal care service at primary health care centers among level of education groups

Ranks				Kruskal-Wallis H	df	Asymp. Sig.
	What is the highest education you ever attended?	N	Mean Rank			
Patient Satisfaction	Unable to read and write	51	184.00	2.735	6	.841
	Read and write	66	188.58			
	Elementary	170	201.16			
	Middle	59	199.07			
	High	11	164.14			
	Diploma	8	224.13			
	Bachelor's degree	25	194.94			
	Total	390				

No statistically significant difference in women's satisfaction with antenatal care service at primary health care centers among level of education groups.

Table 6: Differences in women's satisfaction with antenatal care service at primary health care centers between antenatal visit number groups

Ranks					Mann-Whitney U	Asymp. Sig. (2-tailed)
	Is this your first antenatal visit at this facility for this pregnancy?	N	Mean Rank	Sum of Ranks		
Patient Satisfaction	Yes	183	187.89	34383.50	17547.500	.202
	No	207	202.23	41861.50		
	Total	390				

The study results reveal that there is no statistically significant difference in women's satisfaction with antenatal care service at primary health care centers between antenatal visit number groups.

Table 7: Differences in women's satisfaction with antenatal care service at primary health care centers among family's monthly income groups

Ranks				Kruskal-Wallis H	df	Asymp. Sig.
	Monthly Income	N	Mean Rank			
Patient Satisfaction	<300.000	121	218.83	13.526	3	.004
	301.000 - 600.000	167	184.00			
	601.000-900.000	52	212.30			
	901.000-1200.000	50	159.99			
	Total	390				

There is a statistically significant difference in women's satisfaction with antenatal care service at primary health care centers among family's monthly income groups (p-value = .004).

Table 8: Differences in women's satisfaction with antenatal care service at primary health care centers between type of pregnancy groups

Ranks					Mann-Whitney U	Asymp. Sig. (2-tailed)
	Type of Pregnancy	N	Mean Rank	Sum of Ranks		
Patient Satisfaction	Planned	318	186.64	59350.50	8629.500	.001
	Not planned	72	234.65	16894.50		
	Total	390				

There is a statistically significant difference in women's satisfaction with antenatal care service at primary health care centers between type of pregnancy groups (p-value = .001).

4. Discussion

Pregnancy is a very important and special event from both social and medical points of view. Maternal satisfaction with antenatal care (ANC) services has been linked to the quality of the provided services and the extent to which specific maternal needs are met [15]. The result illustrated by the frequency and percentage in the presented tables, in terms of age (table 1), the researcher found that, most pregnant women were in the age group of (22-28) years at the time of data collection. Followed by those aged (15-21) years. This result is supported by another study in that most of the study sample is within the age group of (20-30) years old [16]. Regarding marital status, the study findings showed that all women participants are married (100.0%). The results are because most of the women got married at an early age according to the prevailing social customs and traditions of the Iraqi family. The fact that most of the Iraqi people are Muslim, and the Islamic religion encourages early marriage [17]. Like the results of a study conducted in Nigeria it was found that the most of participants, three-hundred-sixty-five (94.3%) were married [18].

In terms of residency, most of the study sample (83.8%) were from urban area; this result is supported by a previous study in Erbil governorate, Iraq (2014) their result indicated that most of the participants were urban area, (75.9%) [19]. Related to the educational level, it has been found that less than half are elementary school graduates (43.6%), compared to those who read and write (16.9%); this result was supported by a study in Iraq regarding educational levels most of the sample (43.6%) primary educational, [20]. Another study which agrees with the present study [21]. Regarding monthly income, more than two-fifths reported that their monthly income ranges between (301.000 – 600) Iraqi dinars per month (Table 4-1). This finding is not supported by a descriptive study conducted in Iraq, which indicates most of the participants had a monthly income of less than 300,000 Iraqi dinars [20].

In the present study, it was authenticated the majority of participants (43.8%) have one pregnancy, (table 2) these result is supported by the present result indicated that the majority of this subjects (42.3 %) were one gravida. The results showed that the ideal period for women between the ages of (21-25) years, early marriage [22].

Regarding the number of antenatal care (ANC) visits, The findings disagree with study in Ethiopia ;Jimma Twon, which show the majority of women attended the first antenatal visit (38.9 %), followed by (27.9%) in the second [23][15][24].

In terms of current gestational age, the findings of the present study show that the majority of the pregnant woman (66.4%) were the second trimester, another study indicated that the majority of these subjects (67.8%) were the second trimester. It is considered critical for pregnancy stages of embryo formation. Pregnant needs special attention to nutrition and health status monitor and follow up the scheduled tetanus vaccine and prevent anemia [25].

Concerning parity, type of pregnancy, the study findings disagree with the study in Ethiopia, which shows that the most of pregnant woman (36.6 %) were multipara [26].

In the present study, there was a somewhat satisfactory level of satisfaction with ANC services (84.9%) related to management and treatment, counseling, and other services (tabl 3). Consistent results were revealed in Al-Amara city, Iraq, where mothers' degree of satisfaction with ANC services was with counseling, at level (85.7%) of satisfaction related to different domains of services, such as registration services, diagnostic and treatment, laboratory tests, vaccines, educational cessation, and communication with providers [27]. The current findings also match those of the study of Mansour S et al. (2012) in Al-Madinah Al-Munawaroh, Saudi Arabia, about women's satisfaction with the quality of antenatal care at primary health care centers. The study indicated that nearly two-thirds (62%) of the participants were satisfied with the overall

aspects of antenatal care services[28]. Moreover, the present finding is consistent with the findings of Emelumadu O et al. (2014). They conducted a study titled Perception of Quality of Maternal Healthcare Services among Women Utilizing Antenatal Services in Selected Primary Health Facilities. This study concluded that most women were satisfied with the care they received at health facilities[29]. Furthermore, the current findings disagree with those of Abebe S et al. (2017), who studied the quality of antenatal care services in public health facilities in Ethiopia. A total of 52.6% of the respondents were satisfied with the service provided. Specifically, almost three-fifths (63.4%) of the present study subjects were unsatisfied with the centers' accessibility [30]. The results of the present study disagreed with the findings of Montasser N et al. (2012). They conducted a study on Egyptian women's satisfaction with and perceptions of antenatal care in Shawa village, Dakahlia governorate, Egypt. Their results revealed that nearly three-quarters (72.2%) of their study participants were not satisfied with the accessibility of ANC s[31]. The present finding disagrees with the study of Edie G et al. (2015) about perceptions of antenatal care services by pregnant women attending governmental health centers in the Buea Health District, Cameroon, where they found that only 20% of their study subjects were satisfied with the centers' infrastructure [32].

Concerning to table(4 to 8). The differences in pregnant women's satisfaction with antenatal care service between the groups of residencies, level of education, number of antenatal visits, and monthly income. The study results reveal that there is no statistically significant difference with antenatal care service between the groups of residencies, level of education, number of antenatal visits, while there is a statistically significant difference with monthly income and type of pregnancy. These results are similar with study conducted in Cameroon in 2014 which, noted no association between clients' level of education and satisfaction [33]. While inconsistent with studies in Vietnam. That urban residents were significantly more satisfied with ANC services.

According to a study by Nnebue, on satisfaction with the quality of maternal health care services in Nigeria, the level of satisfaction was not found to be different among women of different socio-economic groups [34].

Moreover, a satisfied woman is more likely to increase compliance with ANC visits. More recently, Seyoum, found that antenatal care service satisfaction among women having one visit is lower than among women having more than one [35].

5. Conclusion

This study revealed that antenatal care services at primary health care centers negatively predict women's satisfaction.

6. Recommendations

1. The Ministry of Health should provide more support and attention to the antenatal care units.
2. Paying attention to the educational and training courses for the health team, especially the nurses who work in pregnancy care units in order to monitor the health status.

7. Conflict of Interest

Nothing will happen to hurt anybody

REFERENCES

1. Ali SA, Dero AA, Ali SA, Ali GB. Factors affecting the utilization of antenatal care among pregnant women: A literature review [Internet]. Available from: <http://www.alliedacademies.org/pregnancy-and-neonatal-medicine/>
- Mahnoosh HA, Fadhil SN. Assessment of Social Factors Associated with Childbirth Satisfaction among Pregnant Women in Al-Nasiriyah City [Internet]. Vol. 10, *Journal of Research in Medical and Dental Science* 1. 2022. Available from: www.jrmds.in
3. Shalash, Ishraq, and Fatin Mohammed. "Assessment of pregnant women's knowledge and practices concerning prenatal care who attend primary health care centers in Baghdad City." *Iraqi National Journal of Nursing Specialties* 25.3 (2012): 1-10
4. Yazdani NS, Haider KA, Khan A, Jaffar Zaidi SA, Rajani A, Nisar I, et al. Pregnant Women's Experiences with Midwifery-Led Antenatal Care Services in Peri-Urban Communities in Karachi, Pakistan. *Patient Relat Outcome Meas.* 2023 May; Volume 14:127–36.
- Khairi SH Naeem FK. Women's Satisfaction Regarding Care Provided by Nurse-Midwives During Labor in Bint AL-Huda Hospital in AL Al-Nasiriyah City. *Iraqi National Journal of Nursing Specialties*; Vol. 32.
- YassenTaha T. Evaluation of clients' satisfaction toward primary health care centers services at Baghdad city. *Iraqi National J Nurs Specialties* .2017; 30.
- Abdullah B, Abdulwahid H. Reproductive-Related Factors Influencing Pregnant Women Satisfaction Toward Vaccination Services. *Journal of Obstetrics, Gynecology and Cancer Research*. 2023 Jul 7;8(4):404–10.
- Khatri RB, Mengistu TS, Assefa Y. Input, process, and output factors contributing to quality of antenatal care services: a scoping review of evidence. *BMC Pregnancy Childbirth*. 2022 Dec 1;22(1).
9. Khleel HA, Mohammed WK. Evaluation of Pregnancy-related Health Behaviors' Change during Pregnancy for Pregnant Women Attending Abo Ghareeb Primary Health Care Sector. *Iraqi National Journal of Nursing Specialties*. Vol 34.
- Jasim BJ, Khalifa MF. Evaluation of Quality of Primary Health Care Services at Primary Health Care Centers in Baghdad City: A Comparative Study. *Iraqi National Journal of Nursing Specialties*. 2018; Vol 31.
- Sameen EK, Abdul K, Jabbar SA. Evaluation of Women's Knowledge and Personal Preference about The Place of Delivery in Baghdad City. *Iraqi National Journal of Nursing Specialties*. 2017; Vol. 30
- Abed HN, Ali IKA. Assessment of Associated Risk Factors with the Incidence Rate of Abortion Cases among Women at Maternity and Pediatric Hospital in Al-Diwaniyah. *Iraqi National Journal of Nursing Specialties*. 2021; Vol 34.
- Khshain WH, Abdulwahid HS. Evaluation of Pregnant Women Health Promotive Behaviors Related to Perceived Severity. *Bahrain Medical Bulletin*. 2023; Vol 45.
14. Alhaqbani SM, Bawazir AA. Assessment of Pregnant Women's Satisfaction with Model of Care Initiative: Antenatal Care Service at Primary Health Care in Cluster One in Riyadh, Saudi Arabia. *Healthcare (Switzerland)*. 2022 Jan 1;10(1).
- Lakew S, Ankala A, Jemal F. Determinants of client satisfaction to skilled antenatal care services at Southwest of Ethiopia: A cross-sectional facility-based survey. *BMC Pregnancy Childbirth*. 2018 Dec 6;18(1).
- Kebede, Dagmawit Birhanu, et al. "Maternal satisfaction with antenatal care and associated factors among pregnant women in Hossana town." *International journal of reproductive medicine* 2020 (2020)
- Saleh, A.M., Othman, S.M., Ismail, K.H., & Shabila1,P.N.(2022). Exploring Iraqi people's perception about early marriage: a qualitative study. *Bio Med Central Women's Health*, 22(393). Retrieved from <https://doi.org/10.1186/s12905-022-01980-y>
- Maternal Satisfaction and Associated Factors Among Pregnant Women Attended at Antenatal Care Service in Bedessa Health Center, Wolaita Zone, Ethiopia, 2018, Assessment of patient satisfaction and perception of quality of antenatal care services among pregnant women attending antenatal clinics at a secondary health care facility in Anambra State, Nigeria
- Shabila, N.; Ahmed, H.; and Yasin, M.; Women's views and experiences of \ antenatal care in Iraq: a Q methodology study. *BMC pregnancy and childbirth*, 2014;14(1):43.

- Yaser A, Hussein A. Assessment of Antenatal Care Services Among Pregnant Women in Al-Hilla City. *Kufa J Nurs Sci* 2015;5(3):192-200.
- Laishram, J.; Thounaojam, U.; Panmei, J.; Mukhia, P.; and Devi, H.; Knowledge and Practice of Ante-natal Care in an Urban Area. *Indian Medical Gazette*, 2013;34(2):101-102.
- Mansur, A.; Rezaul, K.; and Mahmudul, H.; Quality of Antenatal Care in Primary Health Care Centers of Bangladesh. *Journal of family & reproductive health*, 2014;8(4):175.
- Kamil A, Khorshid E. Maternal perceptions of antenatal care provision at a tertiary level hospital, Riyadh. *Oman Med J*. 2013; 28(1):33–5.
- Asifere W, Tessema M, Tebeje B. Clients' Satisfaction with Health Care Providers' Communication and Associated Factors Among Pregnant Women Attending Antenatal Care in Jimma Town Public Health Facilities, Jimma Zone, Southwest. *Int J Pregnancy Childbirth* 2018;4(5):223-30.
25. Emiru, Amanu Aragaw, G. D. Alene, and Gurmesa Tura Debelew. "Women's satisfaction with the quality of antenatal care services rendered at public health facilities in Northwest Ethiopia: the application of partial proportional odds model." *BMJ open* 10.9 (2020): e037085.
26. Fseha B. Assessment of Mothers Level of Satisfaction with Antenatal Care Services Provided at Alganesh Health Center Shire, North West Tigray, Ethiopia. *Biomed J Scient Tech Res* 2019; 16(1):11798-802.
- Abdulhussein Al-Abedi G. Identification of Pregnant Women's Satisfaction among Antenatal Health Care Services in Primary Health Care Centers at Al-Amara City/Iraq. *Bahrain Medical Bulletin*. 2021; Vol 43.
- Mansour S, 1&3 L, Amin H, Fahmy H, 2&4 E. Sahar Mansour Lamadah and Howaida Amin Hassan Fahmy Elsaba. Women's Satisfaction with the Quality of Antenatal Care at the Primary Health Care Centers in Al-Madinah Al-Menawarh, KSA [Internet]. Vol. 9, *Life Science Journal*. 2012; Vol, 9. Available from: <http://www.lifesciencesite.com><http://www.lifesciencesite.com>.643
- Onyeonoro U, Ukegbu A, Emelumadu O, Ezeama N, Ifeadike C, Okezie O. Perception of quality of maternal healthcare services among women utilizing antenatal services in selected primary health facilities in Anambra State, Southeast Nigeria. *Nigerian Medical Journal*. 2014;55(2):148.
- Alemayehu A. Quality of Antenatal Care Service in Public Health Facilities of Chench District, Gamo Gofa Zone, Southern Ethiopia. *Women's Health*. 2017 Feb 21;4(3).
- Abd El-Hamed Montasser N, Mohamad Helal R, Mohamed Megahed W, Khairat Amin S, Mohamed Saad A, Refaat Ibrahim T, et al. Egyptian Women's Satisfaction and Perception of Antenatal Care [Internet]. *Research Article International Journal of TROPICAL DISEASE & Health*. 2012. Vol. 2, Available from: www.sciencedomain.org
- Edie GEHE, Obinchemti TE, Tamufor EN, Njie MM, Njamen TN, Achidi EA. Perceptions of antenatal care services by pregnant women attending government health centers in the buea health district, Cameroon: A cross sectional study. *Pan African Medical Journal*. 2015 May 21;21.
- Gregory Edie Helle Ekana Edie: Thomas Egbe obinchemti Emmanuel njuma tamufor martinimafany njie theophile nana njamen Erick akum achidi (2014) perception of antenatal care services by pregnant women attending government health centers in buea health district, Cameroon: a cross-sections study . *pan African medical journal*, 2015, 21: 45.
- Nnebue C. C, Ebenebe U. E, Edinma E. D, Iyoke C. A, Obionu C. N, Ilika A. L, (2014) Client's knowledge, perception and satisfaction with quality of maternal healthcare service at primary health care level in Nnewi, Nigeria, *Niger journal clinical practitioner*, 2014, 17(5): 594-601.
- Tesfaye T, Mekonnen H, Negesa L. Maternal antenatal care service satisfaction and factors associated with rural health centers, Bursa District, Sidama Zone, Southern Ethiopia: a cross-sectional study. *J Womens Health Care*. 2017;6(363):4-20.