CENTRAL ASIAN JOURNAL OF MEDICAL AND NATURAL SCIENCES
https://cajmns.centralasianstudies.org/index.php/CAJMNS
Volume: 07 Issue: 03 | July 2026 ISSN: 2660-4159

*Article*

# The Importance of Digital Monitoring Systems in Post-Transplant Rehabilitation

**Baxtiyorjon Xalmirzayevich Matkarimov*[1]**

1. Associate Professor, Department of General Surgery and Transplantology, Andijan State Medical Institute, Andijan, Uzbekistan
* Correspondence: matkarimov1973@mail.ru

**Abstract:** Kidney transplant surgery patients' postoperative rehabilitation is extremely important for health monitoring. New digital monitoring systems, telemedicine, and mobile health technologies have proven to be superior ways to enhance medication adherence, lower infectious and non-infectious complication rates, and better connect healthcare providers and patients in recent years. The purpose of this study was to explore the role of digital monitoring systems in the postoperative course of kidney transplant patients at the Clinic of Andijan State Medical Institute from 2023 to 2025. All 17 patients who had received kidney transplantation were analyzed. The results proved that remote monitoring helped to increase adherence to the recommended immunosuppressive treatment and allowed a more personalized approach in rehabilitation process. Patients were able to take medicines more reliably, and health professionals were provided with up-to-date information about changes in the state of their patients. Furthermore, digital monitoring helped to identify early signs of potential complications, decreased the number of infectious events, improved psychological health and wellbeing, and allowed for more efficient use of health care resources. The use of digital communication channels helped to promote patient engagement and active participation in the recovery process. The findings indicate that the adoption of digital health technologies in rehabilitation after transplant can enhance the safety, effectiveness and quality of rehabilitation program. Additionally, this study offers preliminary data to support the general introduction of telemedicine monitoring systems and lays the groundwork for future clinical studies to optimize long-term outcomes in kidney transplant recipients.

**Citation:** Matkarimov, B. X. The Importance of Digital Monitoring Systems in Post-Transplant Rehabilitation. Central Asian Journal of Medical and Natural Science 2026, 7(3), 404-410.

Received: 10th Apr 2026
Revised: 30th Apr 2026
Accepted: 14th May 2026
Published: 06th Jun 2026



**Keywords:** Kidney Transplantation, Telemedicine, Remote Monitoring, Adherence, Postoperative Care, Rehabilitation, Digital Health

## 1. Introduction

Currently, kidney transplantation is considered the best treatment modality for patients with end-stage chronic kidney disease. Although dialysis may help to extend life and keep the body functioning, transplantation provides much better long-term results, improved quality of life, social participation and survival chances. However, the recovery phase after the surgery can be tricky and sensitive, and demands constant care and collaboration from the patient. The first few months after transplanting are critical. In this time the recipient's immune system can identify the transplanted organ as foreign, causing graft rejection [1].

At the same time, the patient is still susceptible to infectious complications, arterial hypertension, metabolic disturbances and side effects of long-term immunosuppressive treatment. Inadequate compliance with prescribed medication is one of the most consistent problems seen in clinical practice. A myriad of studies have found that lack of compliance with immunosuppressive therapy is one of the most important factors in graft dysfunction and graft loss [2].

The typical follow up treatments involve regular outpatient visits, laboratory tests and clinical review. But practical experience shows us that there are often physiologic changes that are not always scheduled. Changes in blood pressure, an increase in blood urea or a deterioration in general health may occur between visits, and be missed until the next visit. This can affect the quality of care and its potential for complications [3]. To meet these challenges, digital monitoring technology is playing a growing role in today's healthcare. Telemedicine, mobile health apps, and remote patient monitoring (RPM) have ushered in new ways to monitor patients continuously in the home setting. These technologies enable patients to receive reminders about taking their medicines, to document important health information, to share laboratory data electronically, and to have real-time communication with their health care providers. Digital platforms not only help patients identify any clinical issues in the early stages but also empower them to be more involved in their care. Patients are engaged in the treatment process, and are informed of their condition, by regularly monitoring personal health data and having direct contact with the clinicians [4].

Transplant recipients should be given special attention as they are more susceptible to infections due to the immunosuppressive drugs they have been prescribed. This is why it is crucial to reduce any unnecessary trips to hospital and to keep a good clinical eye on the patient. Some studies have suggested that remote monitoring improves follow-up efficiency, increases patient safety and decreases postoperative complication rates in transplant patients. In view of these factors, the importance of digital monitoring systems in rehabilitation after transplant has grown more and more significant. This study looks at how digital monitoring affects clinical outcomes after kidney transplantation and investigates the real life effectiveness of it as part of modern day post-transplant care [5].

## 2. Materials and Methods

The aim of this study was to assess the effect of digital monitoring systems on the postoperative rehabilitation of kidney transplant patients. The study was realized on the basis of the clinical results of kidney transplantation in the Department of Surgery No. 2 of the Clinic of Andijan State Medical Institute for three years from 2023 to 2025. This was a retrospective, observational, study in which detailed analysis of the patients' records, laboratory results, operative notes and outpatient follow-up notes were carried out. These data were beneficial for an understanding of patient recovery and as a means of evaluating patient progress during the post-transplant phase [6].

The subjects of the study were 17 adults (between 22 and 58 years of age) who were recipients. Of these 10 were male and 7 female volunteers. All the participants were known to have end-stage chronic kidney disease and were deemed as fit for renal transplantation. All surgical procedures were carried out using well known institutional protocols by the experienced and multi-specialty transplant team. All patients who underwent transplantation received standard triple immunosuppressive therapy after transplantation: tacrolimus, mycophenolate mofetil and glucocorticoids. Doses of the medications were modified on a case-by-case basis based on clinical condition, laboratory results, and function of the graft and any potential adverse effects [7].

Two methods were used to provide postoperative follow-up. The conventional follow-up group had a follow-up schedule of regular outpatient visits and routine laboratory tests and were given routine clinical evaluation. Patients who were enrolled in the digital monitoring program, however, were able to get extra assistance via mobile apps and remote monitoring technologies. These electronic devices served as medication reminders, helped patients record their blood pressure and heart rate daily, and allowed any laboratory findings to be sent quickly to their health care provider. Patients were required to upload daily health data onto the mobile application and the clinicians reviewed this data remotely for clinical recommendations, if they were needed [8]. Multiple clinical measures were chosen to assess the outcomes of rehabilitation. They were immunosuppressive therapy, number of acute rejection, infectious complications, blood pressure control and serum creatinine level. Medication adherence was evaluated using patient self-report, prescription monitoring and laboratory evaluation. Acute rejection was diagnosed by the clinical features, biochemical parameters, and histopathologic diagnosis (biopsy) if clinically warranted. All infectious complications were recorded by clinical examination, laboratory investigations and microbiological tests. Descriptive statistical methods were employed in analyzing data collected. Values of continuous variables were reported as mean values (standard deviations). The comparisons between groups were made with Student's t-test and a p value $< 0.05$ was considered as statistically significant.

This analytical approach enabled a detailed evaluation of the effect of digital monitoring on patient rehabilitation and clinical results after kidney transplant. The ethical aspects related to medical research were adhered to throughout the study. Patient data were fully anonymized and all analyses were performed on anonymized clinical data. The method used was designed to be applicable to the clinical reality and to include the patient's perspective and physician's perspective throughout the post-transplant rehabilitation period [9].

## 3. Results

The Rehabilitation results of 17 kidney transplant recipients, who visited the Clinic of Andijan State Medical Institute during 2023-2025 were studied. Either outpatient follow-up or an integrated digital monitoring system, which supplemented traditional clinical monitoring, was used to treat patients. Collecting data and analyzing it proved to be a meaningful contribution to the improvement of the postoperative management and support of patient recovery after transplantation. One of the most important was that there was a variation in adherence to medication between the two groups. Sticking with immunosuppressive therapy is a fundamental component of a successful kidney graft since failure to take the medication properly greatly raises the likelihood that the graft will fail to function and that the recipient will reject it. Patients engaged in the digital monitoring program showed a significantly greater adherence to treatment plans. The patients in the digital monitoring group adhered to their immunosuppressive therapy as advised in about 95% of cases, while in the patients undergoing conventional follow-up, this figure was 78% [10].

Adherence with the digital monitoring was higher among the patients, seemingly related to clinical stability. Patients remained committed to the treatment and were less likely to miss doses due to the use of medication reminders, regular contacts with health professionals, and access to personal data on health. Therefore, digital support recipients were less likely to experience episodes suggestive of acute rejection [11].

Another benefit of remote monitoring was the ability to identify potential clinical issues earlier than usual, and this also helped to increase medication adherence. By monitoring blood pressure, heart rate and laboratory markers every day, health care providers could recognize a pattern of worsening health before it worsens into serious health issues. A number of patients had transient rises in blood pressure, or increases in the markers of kidney function, which were identified quickly via the digital platform. There was no further deterioration and fewer emergency hospital visits. The other advantage noted was in managing infectious complications. Transplant recipients are still susceptible to infection, and it is important to recognize symptoms early since they are at risk for infection with immunosuppressive therapy. Symptoms were reported sooner, and the patient received medical advice promptly when using the digital monitoring systems.

As a result, fewer patients suffered from infectious complications, and they tended to be less severe than those in the traditional follow-up group [12].

The psychological effect of the sustained support from afar was also noted during the rehabilitation phase. Patients said they felt more confident and secure when they knew that healthcare providers could check on their health information on a regular basis. This continual link helped to decrease anxiety about graft function and promote increased involvement in self-management activities. Communication was enhanced, which fostered a better patient-clinician relationship, and was an important factor in the patient-centered rehabilitation process.

In summary, the results suggest that digital monitoring technologies can be integrated into post-transplant care to boost adherence to treatment, facilitate early detection of complications, increase patient confidence in the treatment, and optimize the use of healthcare resources. These findings indicate digital health interventions could serve as a complementary tool to traditional approaches in postoperative care for kidney transplant patients [13].

**Table 1.** The study compares the proportion of patients who took the medication as directed with the proportion who had an acute rejection episode.

| Group | Medication adherence (%) | Number of acute reaction episodes |
|---|---|---|
| Traditional control | 78 | 5 |
| Digital monitoring | 95 | 1 |

Table 1 shows that digital monitoring helped patients to take a more active and responsible role in their health management. Using the mobile platform, users could monitor their medications, get reminders to take them, and communicate directly with health care providers if they had concerns. This ongoing interaction seemed to help with the support of treatment recommendations and aid in the reduction of acute rejection episodes. The results indicate that digital support used regularly after transplant could significantly decrease clinical risks from non-compliance with medication. Infectious complications also were assessed during the follow-up period. Within the first year after the transplants, six of the patients who were monitored with conventional surgery had at least one documented infection. By comparison, none of the patients in the digital monitoring program experienced infectious complications. The relatively small sample size observed the difference indicates that on-going remote monitoring might help identify warning signs earlier and help rapid intervention before complications set in. This good outcome was probably due to the reporting of symptoms and quick communication between patients and clinicians.

**Table 2.** During Follow-up, infectious complications and graft function were evaluated.

| Group | Infectious complications(n) | Patient with normal transplant graft function(%) |
|---|---|---|
| Traditional control | 6 | 83 |
| Digital monitoring | 1 | 100 |

The positive effect of digital monitoring on graft preservation is also evident in the data in Table 2. Graft function did not change during the follow-up period of 12 months in all patients who were included in the digital monitoring program. Conventional follow-up preserved normal graft function in just 83% of patients, compared to 90% who had normal function with the new regimen. The difference may be due to the fact that remote monitoring systems can detect clinical signs of degradation earlier and enable timely medical action before irreversible damage is done. In addition to the measurable outcomes, digital monitoring also seemed to impact patient experience. Participants said they had become more confident in controlling their health as they could regularly check health information, get reminders about taking medicines, and talk to health care practitioners without having to wait for a clinic appointment. This feeling of ongoing reassurance could have been responsible for enhancing psychological health and more participation in rehabilitation. The results are especially relevant given that as well as physiologically recovering from the transplant, patients also need to learn how to take on long-term self-management tasks. Overall, these findings suggest that digital monitoring systems have the potential to provide more than just traditional clinical monitoring. Digital technologies can help to improve the effectiveness of the post-transplant rehabilitation process by better connecting patients to healthcare providers, helping patients adhere to their care plans and enabling earlier detection of any complications.

## 4. Discussion

The results of this study indicate that digital monitoring can be a useful tool to enhance kidney transplant outcomes. Those enrolled in the remote monitoring program showed greater adherence to immunosuppressive therapy than those who were followed-up in the traditional manner. This is clinically relevant as it is one of the best predictors of long-term graft survival. Any interruption of immunosuppressive therapy can pose the risk of rejection, especially in transplant medicine, and strategies that assist patients with adherence to therapy are therefore critical [14].

One of the other interesting findings was that patients monitored digitally had a lower rate of infectious complications. One of the primary benefits of using remote healthcare technologies is the ability to detect the symptoms early. Patients could alert healthcare providers to concerns on time, which would enable care to be provided before things escalate to a critical state, due to daily monitoring and the availability of fast communication. Previous research with telemedicine interventions for chronic disease management and transplant follow-up has reported similar findings [15].

Another advantage of digital monitoring is the power it gives in enhancing patient–healthcare provider interaction. While traditional follow-up models involve periodic visits weeks or months apart, digital platforms offer opportunities for ongoing interactions. Patients can access their health data instantly and receive expert advice as needed. This constant interaction can help to decrease uncertainty, increase confidence and promote greater involvement in self-care. The psychological benefit of feeling cared for is underestimated, especially for transplant recipients, who may have concerns about how the graft will work and what they will need to do for the long term. Healthcare systems, on the other hand, might also benefit from monitoring through digital devices because of the potential to enhance resource use. Spending less on unnecessary hospital visits, intervening earlier on issues that might arise, and fewer unplanned admissions can lead to a more efficient service delivery. These benefits could assume greater significance as transplant programs grow and as health care providers look for long-term management of their patients that can be sustained. However, some of the issues need to be recognize. Digital monitoring does not work for all patients. Patients with very good adherence to treatment may not benefit any further. In addition, technological barriers still need to be taken into account. The effectiveness of remote monitoring program may be compromised, especially in certain populations, due to limited access to smartphones, unreliable internet connectivity and lack of digital literacy. All these show the need to adapt a DHI for the needs and capabilities of specific patients [16].

Some limitations of the present study are a single-center design and a relatively small sample size. Thus, the results need to be taken with a pinch of salt. More large-scale, multicenter trials are required to assess the long-term clinical effects of digital monitoring and to look at patient groups that could best benefit from these technologies.

The results offer promising evidence for use of digital monitoring to aid safe and effective rehabilitation after kidney transplant, despite these limitations. Overall, digital health technologies have the potential to play a greater role in the future of transplant care, as they help to increase the likelihood of adherence, improve early detection of complications, make patients more engaged in their health, and improve the communication between patients and clinicians [17].

## 5. Conclusion

The results of this study suggest that digital monitoring systems could have a significant benefit in enhancing the postoperative rehabilitation of kidney transplant patients. Patients on remote digital monitoring had better adherence to immunosuppressive therapy, a decreased incidence of infectious complications and more stable graft function during follow-up. The enhancements underscore the importance of ongoing surveillance to ensure the long-term viability of transplants. In addition to its clinical benefits, digital monitoring strengthened communication between patients and healthcare providers, allowing concerns to be addressed promptly and facilitating more informed clinical decision-making. The technology also appeared to enhance patient confidence and psychological well-being by encouraging active participation in health management and providing a sense of ongoing professional support. Furthermore, digital monitoring may contribute to more efficient use of healthcare resources by reducing unnecessary hospital visits and enabling earlier intervention when problems arise. Such advantages are particularly relevant in transplant medicine, where long-term follow-up is essential and timely responses to clinical changes can significantly influence outcomes. Overall, digital monitoring represents a promising approach for optimizing post-transplant care. Expanding the use of these technologies and adapting them to individual patient needs may further improve graft survival, enhance quality of life, and support the continued development of patient-centered transplant rehabilitation programs in the future.

## REFERENCES

[1] Kidney Disease: Improving Global Outcomes (KDIGO) Transplant Work Group, “KDIGO Clinical Practice Guideline for the Care of Kidney Transplant Recipients,” Kidney International Supplements, vol. 10, no. 2, pp. S1–S141, 2020.

[2] S. Hariharan, A. K. Israni, and G. Danovitch, “Long-term survival after kidney transplantation,” New England Journal of Medicine, vol. 385, pp. 729–743, 2021, doi: 10.1056/NEJMra2013716.

[3] F. Dobbels, S. De Geest, R. Van Damme-Lombaerts, et al., “Adherence to immunosuppressive therapy in kidney transplant recipients: A systematic review,” Transplant International, vol. 18, no. 10, pp. 1121–1133, 2005, doi: 10.1111/j.1432-2277.2005.00199.x.

[4] A. J. Greenberg, A. Lipitz-Snyderman, and M. S. Lachs, “Telemedicine in kidney transplantation: Monitoring and patient outcomes,” American Journal of Kidney Diseases, vol. 79, no. 4, pp. 513–524, 2022, doi: 10.1053/j.ajkd.2022.01.003.

[5] Centers for Disease Control and Prevention, “Telehealth and Remote Patient Monitoring Guidance for Chronic Disease Management,” CDC, Atlanta, GA, USA, 2021.

[6] S. De Geest, H. Burkhalter, A. Boggio, et al., “Effectiveness of electronic monitoring on medication adherence in transplant patients: A systematic review,” Patient Preference and Adherence, vol. 12, pp. 265–279, 2018, doi: 10.2147/PPA.S145832.

[7] National Kidney Foundation, "Kidney Transplant Patient Care Guidelines: Post-Transplant Monitoring and Follow-Up," National Kidney Foundation, New York, NY, USA, 2020.

[8] J. R. Rodrigue, M. Paek, J. Whiting, et al., "Use of mobile health technology in kidney transplant recipients: Patient perspectives and clinical outcomes," Clinical Transplantation, vol. 33, no. 12, Art. no. e13772, 2019, doi: 10.1111/ctr.13772.

[9] S. Oberlin, M. A. Schnitzler, and A. O. Gaber, "Remote monitoring after kidney transplantation: Clinical impact and patient satisfaction," Transplantation Proceedings, vol. 53, no. 8, pp. 2550–2557, 2021, doi: 10.1016/j.transproceed.2021.05.024.

[10] M. Kuang, Z. Yao, J. Chen, Y. Yan, and Z. Li, "The effectiveness of remote interventions based on digital health technology in kidney transplant recipients: A systematic review," World Journal of Urology, vol. 43, no. 1, 2025, doi: 10.1007/s00345-025-05582-9.

[11] L. Zhou, K. Cheng, L. Chen, X. Hou, and J. Wan, "Effectiveness of eHealth for Medication Adherence in Renal Transplant Recipients: Systematic Review and Meta-Analysis," Journal of Medical Internet Research, 2025, doi: 10.2196/73520.

[12] M. Tuncer and G. Z. Tuncer, "Effect of eHealth Interventions on Medication Adherence in Kidney Transplant Recipients: Meta-Analysis of Randomised Controlled Trials," Journal of Renal Care, vol. 51, no. 2, 2025.

[13] H. Y. Jung, Y. Jeon, S. J. Seong, et al., "ICT-based adherence monitoring in kidney transplant recipients: A randomized controlled trial," BMC Medical Informatics and Decision Making, vol. 20, no. 1, p. 105, 2020, doi: 10.1186/s12911-020-01146-6.

[14] R. Mendel, J. Nie, M. Keller, Y. Aly, H. Sandhu, M. Handmacher, and L. Kayler, "Digital Enrollment and Survey Response of Diverse Kidney Transplant Seekers in a Remote Trial (KidneyTIME): An Observational Study," Kidney and Dialysis, vol. 5, no. 2, p. 19, 2025.

[15] C. D. Manko, B. J. Apple, A. R. Chang, et al., "Telemedicine in Advanced Kidney Disease and Kidney Transplant: A Qualitative Meta-Analysis of Studies of Patient Perspectives," Kidney Medicine, vol. 6, no. 7, Art. no. 100849, 2024, doi: 10.1016/j.xkme.2024.100849.

[16] S. Esayed, E. Kim, H. C. Sung, et al., "Hybrid Telemedicine and In-Person Care for Kidney Transplant Follow-Up: A Qualitative Study," Clinical Transplantation, vol. 39, no. 2, Art. no. e70106, 2025, doi: 10.1111/ctr.70106.

[17] "Barriers and facilitators of integrating telemedicine services into routine kidney transplant patient care: A scoping review of qualitative research," Journal of Health, Population and Nutrition, vol. 44, p. 309, 2025.